

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 July 2002 (18.07.2002)

PCT

(10) International Publication Number
WO 02/055026 A2

- (51) International Patent Classification⁷: A61K
- (21) International Application Number: PCT/US02/00839
- (22) International Filing Date: 11 January 2002 (11.01.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/261,525 12 January 2001 (12.01.2001) US
- (71) Applicant (for all designated States except US): AFFY-MAX, INC. [US/US]; 4001 Miranda Avenue, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HOLMES, Christopher [US/US]; 13633 Westover Drive, Saratoga, CA 95070 (US). PAN, Yijun [US/US]; 3791 Boswell Terrace, Fremont, CA 94536 (US).

(74) Agents: HOSTETLER, Michael, J. et al.; Brobeck, Phleger & Harrison, LLP, 12390 El Camino Real, San Diego, CA 92130 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/055026 A2

(54) Title: PERFLUORO SULFONYL HALIDES AND RELATED SPECIES AS POLYMER SUPPORT MODIFIERS



(I)

(57) Abstract: Activated Supports, support-bound activators, strongly acidic supports, and silylating supports are provided; the activated support having the formula (I) wherein L is a linking group component; X is F, Cl, OH, and trisubstituted silyloxy; and the shaded circle represents a solid or semi-solid support. Methods of using the activated supports in solid phase organic synthesis are also provided.

**PERFLUORO SULFONYL HALIDES AND RELATED SPECIES
AS POLYMER SUPPORT MODIFIERS**

CROSS-REFERENCES TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/261,525
5 filed January 12, 2001.

**STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER
FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT**

Not applicable

BACKGROUND OF THE INVENTION

10 The present invention relates to the field of linking groups or activators that are useful in the solid-phase preparation of single compounds and libraries, as well as methods that employ such linking groups and activators. The present invention also relates generally to the use of activated supports and their use in organic solution phase chemistry and as derivatizing agents to aid in chromatography.

15 Solid phase synthesis has attracted considerable attention from the scientific community, and in particular, the pharmaceutical and agricultural research communities in an effort to speed up the discovery of new biologically active compounds. Critical to the solid phase synthesis of such compounds are the means to attach (and subsequently remove) the compounds from a support. The most frequently used linkers are acid-labile
20 and photo-labile linkers which typically result in the cleaved product having a residual functional group (e.g., carboxylic acid, amide, amine or hydroxy group). Recently, the term "traceless linker" has been used to describe a strategy of releasing compounds from a solid support with little or no trace of the original point of attachment. See James, *Tetrahedron Lett.*, 1999, 55, 4855; Andres, *et al.*, *Curr. Opin. Chem. Biol.*, 1998, 2, 353;
25 Reitz, *Curr. Opin. Drug Discovery Dev.*, 1999, 2, 358; and Zaragoza, *Angew. Chem., Int. Ed.* 2000, 39, 2077.

What is needed in the art are polymer support-linker species that activate certain molecules toward other transformations that would be useful in the preparation of single compounds or compound libraries, and that can also act as traceless linkers.

NAFION™ (Dupont, Wilmington, Delaware) is a perfluororesinsulfonic acid that could, in principle be used as a traceless linker in solid-phase organic synthesis. However, the inability to utilize NAFION™ resin in high yields and conversions in solid phase organic synthesis applications has been noted. See Akhtar, *et al.*, *Tetrahedron Lett.* 2000, 41, 4487; Liu, *et al.*, *Tetrahedron Lett.* 2000, 41, 4493. Such limitations include the inability to be wetted or swollen by most aprotic organic solvents. Thus, although NAFION™ is robust and chemically resilient, it is not useful in solid-phase organic synthesis because the perfluoropolymer side chains are not solvated. As a result, there is a need in the art for a polymer-supported perfluorosulfonate that is swellable and wettable by most common solvents.

In addition to being used for solid phase organic synthesis, solid and semi-solid supports are also used in solution phase organic chemistry as catalysts or reagents. For example, in the field of catalysis and organic chemistry, there is widespread use of highly acidic catalysts to promote chemical transformations. These often take the form of polymer supported acids, which can be readily filtered away from a reaction mixture at the completion of the reaction. Resins known to those skilled in the art include polystyrene-based resins, controlled pore glass beads, NAFION™, polyethylene glycol resins, TENTAGEL™ (Rapp Polymere GmbH, Tübingen, Germany), and the like. See Olah, *Synthesis* 1986, 7, 513. Whereas ion-exchange resins, and in particular cation exchange resins, are polymer-supported acids, they are typically based on phenylsulfonic acids and are hence limited in acidity. Perfluorosulfonic acids are dramatically more acidic than phenylsulfonic acids and as such, are quite distinct and often capable of catalyzing a broader range of chemical transformations. Thus, there is a general need in the art for polymer-supported perfluorosulfonic acids and related derivatives.

Processes for making perfluorosulfonic solid acids via encapsulation of perfluorsulfonic acids into hydrocarbon resins have been disclosed in WO 98/30521, however, these are not expected to find application in the field of solid phase organic synthesis because the perfluorosulfonate groups are not covalently bound to the resin.

NAFION™ has been demonstrated to act as a highly acidic supported catalyst. Its properties have been noted by Olah, *Synthesis* 1986, 7, 513; Yamoto, *Recent Res. Devel. In Pure & Applied Chem.* 1998, 2, 297; Harmer, *Adv. Mater.* 1998, 10, 1255. However, NAFION™ and related polymers in the art are not effectively swollen by most aprotic


organic solvents. Because NAFION™ is not swellable or wettable by most common organic solvents, only the acid groups on the surface of NAFION™ are available for reaction, while the majority of the acid groups contained within the polymer are unavailable for reaction. One technique of increasing the effective surface area is to grind
5 the polymer into fine particles and to imbed these into an inert carrier such as clay or amorphous silica. See Harmer, *Adv. Mater.* 1998, 10, 1255. These hybrid materials are subject to leaching artifacts and can often be difficult to filter away due to the heterogeneity of the particles. Thus, there is a need in the art for supported superacids that are swollen by common organic solvents, allowing supported superacids to be used for
10 more applications and under milder reaction conditions. See Ishihara, *et al.*, *Angew. Chem. Int. Ed.* 2001, 40.

A superacid polymer-supported resin that is swellable would be highly desirable due to the ease of use, the enhanced effective acid content (because most or all of the contained acid groups would be available for use), and compatibility with a wide range of
15 solvents.

SUMMARY OF THE INVENTION


The present invention provides a variety of support-bound moieties and activated supports, that have utility in solid phase or solution phase synthesis, biosynthesis,
20 catalysis, purification, analysis, and identification and screening. Each moiety and activated support includes an activator portion that serves as a reactive center and a linking group component that serves to provide a robust linkage between the support and the activator portion. The linking group components further include an activator enhancing portion that serves to increase the reactivity of the activator portion and suitable spacer
25 that provides sufficient distance between the activator portion and the support.

In one aspect, the present invention provides a support-bound activator having the formula:




wherein L is a linking group component; X is a member selected from the group consisting of F, Cl, OH, and trisubstituted silyloxy; wherein the support-bound activator is covalently attached to a solid or semi-solid support.

In another aspect, the present invention provides an activated support comprising a solid or semi-solid support; and at least one support-bound activator having the formula:




wherein L is a linking group component; X is a member selected from the group consisting of F, Cl, OH, and trisubstituted silyloxy; and wherein the support-bound activator is covalently attached to the solid or semi-solid support.

In another aspect, the present invention provides a support-activated target comprising a solid or semi-solid support; an activating group covalently attached to the solid or semi-solid support, wherein the activating group has the formula:




wherein L is an linking group component; and a target group covalently attached to the solid or semi-solid support; wherein the target group can be cleaved from the activating group by a nucleophile.

In another aspect, the present invention provides a library of support-activated targets comprising a plurality of support-activated target members, wherein each support-activated target member further comprises a solid or semi-solid support; an activating group covalently attached to the solid or semi-solid support, wherein the activating group has the formula:




wherein L is an linking group component; and a target group covalently attached to the activating group; wherein the target group of at least one support-activated target member in the library is different from the target group of at least one other support-activated target member in the library.

- 5 In another aspect, the present invention provides a strongly acidic support comprising a solid or semi-solid support; and at least one support-bound strongly acid group having the formula:



- 10 wherein L is a linking group component; and X is OH; and wherein the support-bound strongly acid group is covalently attached to the solid or semi-solid support.

In another aspect, the present invention provides a silylating support comprising a solid or semi-solid support; and at least one support-bound silylating group having the formula:




- 15 wherein L is a linking group component; X is a trisubstituted silyloxy; wherein the support-bound silylating group is covalently attached to the solid or semi-solid support.

In yet another aspect, the present invention provides a method for covalently attaching a nucleophile to a compound having a hydroxy group or an enolizable ketone, the method comprising,

- 20 (a) contacting a compound having a hydroxy group or an enolizable ketone with a support-bound activator, wherein said contacting a compound having a hydroxy group or an enolizable ketone with a support bound activator forms an activated complex; and
 (b) contacting the activated complex with a reagent comprising a nucleophile under conditions sufficient to covalently attach the nucleophile to the compound.

In still another aspect, the present invention provides a linker reagent having the formula:



wherein L is a linking group component; and A is an attaching group.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a synthetic scheme for the preparation of a support-bound perfluorosulfonyl fluoride linker in accordance with the present invention.

Figure 2 illustrates a synthetic scheme for the preparation of a support-bound perfluorosulfonic acid in accordance with the present invention.

10

Figure 3 illustrates a synthetic scheme for the deoxygenation reaction sequence for phenols to generate the parent arenes in accordance with the present invention.

Figure 4 illustrates a synthetic scheme to effect a cleavage/cross coupling reaction to afford biaryls in accordance with the present invention

15

Figure 5 illustrates a synthetic scheme whereby a library of targets are prepared on the polymer support in accordance with the present invention.

Figure 6 illustrates the synthesis of a known drug substance and subsequent cleavage from the polymer support in accordance with the present invention.

Figure 7 illustrates the preparation of a perfluorosulfonic acid polymer in accordance with the present invention.

20

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention and as used herein, the following terms and abbreviations are defined with the following meanings, unless explicitly stated otherwise. These explanations are intended to be exemplary only. They are not intended to limit the terms as they are described or referred to throughout the specification. Rather these explanations are meant to include any additional aspects and/or examples of the terms as described and claimed herein.

25

The following abbreviations are used herein: AcOH or HOAc, acetic acid; Boc, t-butoxycarbonyl; DMF, dimethylformamide; EtOAc, ethyl acetate; NMP, N-methylpyrrolidone; TFA, trifluoroacetic acid.

The term "support-bound activator" as used herein refers to an activator which is
5 attached to a solid or semi-solid support by at least one covalent bond. The term
"activator" as used herein refers to a chemical moiety which activates, or increases the
chemical reactivity of, another portion of a molecule. The term "support-bound activator"
is not meant to limit the location of the support-bound activator on the solid or semi-solid
support; that is, the support-bound activator can be covalently attached to the surface
10 portion of the solid or semi-solid support or the support-bound activator can be covalently
attached to any internal portion of the solid or semi-solid support. In other words, the
support-bound activator can be covalently attached to any portion of the solid or semi-
solid support.

The term "activator portion" as used herein refers to the portion of an activator that
15 increases the chemical reactivity of the activator. In a preferred embodiment, an activator
portion comprises the moiety $(-\text{CF}_2-\text{SO}_2-\text{X})$. Activator portions may have analogous
counterparts known in the art of solution phase chemistry.

The term "linking group component" as used herein means a moiety that links
together a plurality of other moieties. Thus, in one embodiment of the present invention,
20 one portion of a linking group component in a surface-bound activator will have at least
one covalent bond with the solid or semi-solid support, and another portion of the linking
group component will have at least one covalent bond with an activator portion.
Furthermore, the linking group component of a surface-bound activator may have a
covalent bond with a plurality of activator portions. Preferably, the linking group
25 component provides suitable spacing for the activator portion to interact with molecules
exposed to the activator. The linking group component is preferably 6-50 atoms long,
more preferably 8-40 atoms long, even more preferably 8-30 atoms long, and yet more
preferably 8-20 atoms long. Additionally, the linker reagent, prior to reaction with the
solid or semi-solid support, will include a linking group component; this linking group
30 component will have one portion that has at least one covalent bond with an attaching
group and another portion with at least one covalent bond with an activator portion. The

term "attaching group" as used herein refers to that portion of a linker reagent that can form a covalent bond with the solid or semi-solid support.

The term "activator enhancing portion" as used herein refers to that portion of the linking group component that increases the chemical reactivity of the activator portion of a support-bound activator. An activator enhancing portion may also extend the length of the linker group component, and additionally, effect the motility of the surface-bound activator.

The terms "solid" or "semi-solid support" as used herein refers to any form of a polymer or composite material that does not completely dissolve in a solvent. By way of example only, a solid or semi-solid support includes colloids (isolated or in suspension), gels, resins, films, as well as any other form of a polymer or composite materials that retains a distinct identity apart from the solvent. The term is not meant to limit in any manner the size, shape, form, or chemical structure of the polymeric or composite material. Such polymeric or composite materials are well known in the art, including, by way of example only, cellulose, pore-glass, silica, polystyrene, polystyrene cross-linked with divinylbenzene, polyacrylamide, latex, dimethylacrylamide, dimethylacrylamide cross-linked with N,N'-bis-acryloyl ethylene diamine, glass, glass coated with a hydrophobic polymer, composites, or any other material conventionally used in solid phase organic synthesis. A valuable reference in this regard is the Novabiochem 2000 catalog, which is incorporated herein by reference. In addition, the term solid or semi-solid support is not limited by the presence and nature of cross-linking groups, and by the nature of the exposed functional groups. Exposed functional groups are moieties on the solid or semi-solid support that can react with linker reagents to form support-bound activators; preferred exposed functional groups include -OH, -SH, -NH₂, silyloxy, NHR, NH₂NH, CO₂H, CO₂R, C(O)H, -Br, -I, halomethyl, and alkenyl. Furthermore, the exposed functional groups can be located on the surface of the solid or semi-solid support or dispersed throughout the solid or semi-solid support. In one embodiment, the solid or semi-solid support has a rigid or semi-rigid surface.

The term "particulate support" as used herein refers to a type of solid or semi-solid support that is in the form of small particles. The term is not meant to limit the shape of the solid or semi-solid support. Thus, by way of example only, a particulate support can be a sphere, disk, pellet, sheet, plug, pin, crown, lantern, capillary, hollow fiber, needle, solid

fiber, in a beaded or non-beaded form, a resin, a gel, a microsphere, an amorphous shape, or any other conventional form. As one skilled in the art will readily recognize, the scope of the present invention is not limited to the form or shape of the particulate support. The term "particulate support" is not meant to limit the chemical structure of the solid or semi-
5 solid support, and can be composed of any polymer, composite, cross-linker, if any, and exposed functional group. Thus, by way of example only, a particulate support can be composed of cellulose, pore-glass, silica, polystyrene, polystyrene cross-linked with divinylbenzene, polyacrylamide, latex, dimethylacrylamide, dimethylacrylamide cross-linked with N,N'-bis-acryloyl ethylene diamine, glass, glass coated with a hydrophobic
10 polymer, composites, or any other material conventionally used in solid phase organic synthesis. A valuable reference in this regard is the Novabiochem 2000 catalog, which is incorporated herein by reference. A particulate support may also be porous, deformable, hard, wettable, or swellable. The particles will generally be at least 20 micron, preferably at least 75 micron, and more preferably at least 100 micron in diameter. A particulate
15 support will generally maintain its mechanical integrity during use, have functional groups that can react with active species, allow for the serial synthesis of attached targets, can be readily mixed and separated, and will allow for convenient detachment of tags and products. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-
20 through devices.

The term "resin" as used herein refers to any of a class of solid or semi-solid organic products of natural or synthetic origin, generally of high molecular weight with no definite melting point. A resin may be chemically inert toward reagents and solvents used in solid phase syntheses or may be functionalized with reactive moieties. Resins may
25 swell extensively in solvents.

The term "gel" as used herein refers to a colloidal suspension of a liquid in a solid, forming a jellylike material having properties of both solid and a solution.

The term "colloid" as used herein a substance consisting of very tiny particles suspended in a continuous medium, such as a liquid, a solid, or a gaseous substance.
30 Generally, the diameter of a colloidal particle is from 20 nm to 200 micron, more preferably 100 nm to 200 micron, even more preferably 500 nm to 200 micron, and yet even more preferably 1 micron to 200 micron.

The term "microsphere" as used herein refers to any material which is roughly spherical in shape. Microspheres may be processed, machined, milled, ground, or extruded according to processes known in the art.

5 The term "superacid" as used herein is a solution of a strong acid in a very acidic solvent. A superacid is an acid that exhibits strength greater than that of 100% H₂SO₄.

The term "controlled pore glass" as used herein refers to a glass used as an inorganic support. Controlled pore glass is produced from a borosilicate base material which is heated to separate the borates and the silicates. The borates are leached out from the material, leaving the silica glass with uniform, controlled pores. Controlled pore
10 glasses have excellent mechanical properties, and can be prepared with a wide range of porosities and average pore sizes. They can be modified to include a variety of functional groups.

The term "polyacrylamide" or PAM as used herein refers to a synthetic water-soluble polymer made from monomers of acrylamide. PAM may be fashioned into gels
15 having a variety of pore sizes.

The term "polyethylene glycol" or "PEG" as used herein refers to a water-soluble, waxy polymer comprising subunits HO-(CH₂CH₂O)_nH. The term
"poly(ethyleneglycol)monomethyl ether" as used herein refers to CH₃O(CH₂CH₂O)_nH and may be referred to as methyl-capped PEG. A skilled artisan will recognize that other
20 capped PEGs may be employed without departing from the spirit of the invention.

The term "silica gel" as used herein refers to silicic acid or precipitated silica. Silica gel is an amorphous powder insoluble in water and organic solvents. Silica gel can adsorb up to about 40% of its weight in moisture.

The term "cellulose" as used herein refers to one of many polymers found in
25 nature, as for example in wood, paper, and cotton. Cellulose is a polysaccharide and comprises repeating units of the monomer glucose. Cellulose may be crosslinked and derivatized, e.g., as in methylcellulose.

The term "acrylic acid grafted polypropylene" as used herein refers to a material comprising a polypropylene backbone and polyacrylate, e.g. acrylic acid side chains. One
30 process modifying the properties of polyolefins comprises the "grafting" of polar monomers onto the polyolefin.

The term "target group" as used herein refers to a moiety covalently bound to a support-bound activator. According to one embodiment of the present invention, a support-bound target comprises a target group covalently attached to support as disclosed herein.

5 The term "cleaved from" as used herein refers to a plurality of transformations disclosed herein in which a target group is removed from an activated support. According to one embodiment of the invention, a support-bound target group is cleaved from or released from a support-bound activator.

10 The term "polystyrene support" as used herein refers to polymerized monomers of styrene. A polystyrene support may take the form of beads as described herein. A polystyrene support may comprise a cross-linked copolymer of styrene and divinyl benzene.

15 The term "polystyrene modified by polyethylene glycol" as used herein refers to a polystyrene polymer or cross-linked polystyrene polymer grafted with ethylene glycol monomer or ethyleneglycol polymer (PEG). An example of a polystyrene modified by polyethylene glycol is TENTAGELTM resin.

 The term "grafted" as used herein refers to a process wherein monomers or polymers are covalently bonded to an existing polymer.

20 The term "TENTAGELTM" as used herein refers to a family of grafted copolymers consisting of a low crosslinked polystyrene matrix on which polyethyleneglycol (PEG) is grafted. Because PEG is a polymer with both hydrophobic and hydrophilic properties, the graft copolymer shows modified physico-chemical properties. A TENTAGELTM resin may comprise about 50 - 70 % PEG (w/w). Therefore, many chemical and physical properties of these polymers are highly dominated by the properties of PEG and not by the

25 hydrophobic polystyrene support. Other related members of this family, for the purpose of this invention, include ARGOGELTM resin (Argonaut Technologies, Foster City, California), ARGOPORETM resin (Argonaut Technologies, Foster City, California), HYPOGELTM resin (Rapp Polymere GmbH, Tubingen, Germany), and JANDAJELTM resin (Scripps Research Institute, La Jolla, California).

30 The term "protecting group" as used herein, refers to any of the groups which are designed to block one reactive site in a molecule or moiety while a chemical reaction is carried out at another reactive site. More particularly, the protecting groups used herein

can be any of those groups described in Greene, et al., Protective Groups In Organic Chemistry, 2nd Ed., John Wiley & Sons, New York, NY, 1991, which is incorporated herein by reference. The proper selection of protecting groups for a particular synthesis will be governed by the overall methods employed in the synthesis and such practice is well known to those skilled in the art. The term "protected forms thereof" as used herein refers to a moiety to which a protecting group has been attached.

The term "unsubstituted" refers to molecules or moieties that do not have additional moieties attached to the named group other than the root compound. Thus, an unsubstituted (C₁-C₈)alkyl group consists only of from 1 to 8 alkyl carbon atoms with attached hydrogens.

The term "optionally substituted" or "substituted" refers to molecules or moieties substituted by one to four substituents, independently selected from lower alkyl, lower aryl, lower aralkyl, lower alicyclic, hydroxy, lower alkoxy, lower aryloxy, perhaloalkoxy, aralkoxy, heteroaryl, heteroaryloxy, heteroarylalkyl, heteroaralkoxy, azido, amino, guanidino, halo, lower alkylthio, oxo, acylalkyl, carboxy esters, carboxyl, carboxamido, nitro, acyloxy, aminoalkyl, alkylaminoaryl, alkylaminoalkyl, alkoxyaryl, arylamino, aralkylamino, phosphono, sulfonyl, carboxamidoalkylaryl, carboxamidoaryl, hydroxyalkyl, haloalkyl, alkylaminoalkylcarboxy, aminocarboxamidoalkyl, cyano, lower alkoxyalkyl, lower perhaloalkyl, and arylalkyloxyalkyl.

The term "lower" referred to herein in connection with organic radicals or compounds respectively defines such as with up to and including 10 carbon atoms, preferably up to and including 6 carbon atoms, and advantageously one to four carbon atoms. Such groups may be straight chain, branched, or cyclic.

The term "alkyl" refers to a saturated hydrocarbon radical which may be straight-chain (for example, methyl, ethyl, propyl or butyl) or branched-chain (for example, isopropyl, t-amyl, or 2,5-dimethylhexyl) or cyclic (for example, cyclobutyl, cyclopropyl or cyclopentyl). This definition applies both when the term is used alone and when it is used as part of a compound term, such as "aralkyl" and similar terms. Preferred alkyl groups are those containing from 1 to 8 carbon atoms, and are referred to as a "(C₁-C₈)alkyl" group. Other preferred alkyl groups include those containing from 1 to 20 carbon atoms. All numerical ranges in this specification and claims are intended to be inclusive of their upper and lower limits. The alkyl group may be optionally substituted.

The term "alkylene" as used herein refers to a divalent carbon moiety, e.g., CR_2 . Alkylene groups be linked to form straight chain alkyl groups and may be optionally substituted with up to two substituents per carbon atom.

5 The term "alkenyl" as used herein refers to a moiety which contains one or more sites of unsaturation. The term "alkenyl" as used herein may also refer to a moiety which contains at least one carbon-carbon double bond and includes straight-chain, branched-chain and cyclic groups. Alkenyl groups may be optionally substituted. The terms "vinyl" and "olefinic" are sometimes used interchangeably with the term alkenyl.

10 The term "($\text{C}_1\text{-C}_8$)alkenyl" as used herein refers to an alkenyl group having from 1 to 8 carbon atoms. Such groups may be straight chain, branched, or cyclic. A ($\text{C}_1\text{-C}_8$)alkenyl group may be optionally substituted.

The term "alkynyl" as used herein refers to a moiety that contains at least one carbon-carbon triple bond and includes straight-chain, branched-chain and cyclic groups. Alkynyl groups may be optionally substituted.

15 The term "($\text{C}_1\text{-C}_8$)alkynyl" as used herein refers to an alkynyl group having from 1 to 8 carbon atoms. Such groups may be straight chain, branched, or cyclic. A ($\text{C}_1\text{-C}_8$)alkynyl group may be optionally substituted.

20 The term "aryl" as used herein refers to a moiety having either a single ring or multiple rings which are fused together and which has at least one ring having a conjugated pi electron system; the multiples rings may also be linked covalently or via a common group such as an ethylene or methylene moiety. Aryl groups may be optionally substituted at any position on the ring which would otherwise be occupied by a hydrogen atom.

25 The term "trisubstituted silyloxy" as used herein refers to a moiety having the formula $-\text{OSiR}^3\text{R}^4\text{R}^5$, wherein, each of R^3 , R^4 , and R^5 is independently a member selected from the group consisting of substituted ($\text{C}_1\text{-C}_8$)alkyl, unsubstituted ($\text{C}_1\text{-C}_8$)alkyl, substituted ($\text{C}_1\text{-C}_8$)alkenyl, unsubstituted ($\text{C}_1\text{-C}_8$)alkenyl, substituted aryl, and unsubstituted aryl.

30 The term "heteroalkyl" refers to an alkyl radical or group in which 1, 2, or 3 of the carbon atoms in the main chain of atoms has been replaced by a heteroatom selected from oxygen, nitrogen, sulfur or silicon. Thus, the term ($\text{C}_2\text{-C}_8$)heteroalkyl refers to a group having from 2 to 8 main chain atoms, at least one of which is a heteroatom. For example,

a C₃ heteroalkyl group is meant to include -CH₂OCH₃ (the oxygen atom taking the place of the central carbon atom in a C₃ alkyl (propyl) group. A heteroalkyl may be optionally substituted.

5 The term "heteroaryl" as used herein refers to an aryl radical having from 1 to 4 heteroatoms as ring atoms in the aromatic ring, with the remainder of the ring atoms being carbon atoms. Suitable heteroatoms include oxygen, sulfur, and nitrogen. By way of example only, suitable heteroaryl groups include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolyl, pyridyl-N-oxide, pyrimidyl, pyrazinyl, and imidazolyl. A heteroaryl may be optionally substituted.

10 The term "biaryl" as used herein refers to an aryl radical containing more than one aromatic ring including both fused ring systems and aryl groups substituted with other aryl groups. By way of example only, suitable biaryls include naphthyl and biphenyl. A biaryl may be optionally substituted.

The term "alicyclic" as used herein refers to a moiety or a compound which
15 combines the structural properties of aliphatic and cyclic or heterocyclic compounds and includes, but is not limited to, aromatic, cycloalkyl and bridged cycloalkyl compounds. An alicyclic may be optionally substituted. By way of example only, suitable alicyclics include cyclohexenylethyl and cyclohexylethyl are suitable alicyclic groups.

The term "aralkyl" refers to an alkyl group substituted with an aryl group. Suitable
20 aralkyl groups include benzyl, picolyl, and the like, and may be optionally substituted.

The terms "arylamino" (a), and "aralkylamino" (b), respectively, refer to the group -NRR' wherein respectively, (a) R is aryl and R' is hydrogen, alkyl, aralkyl or aryl, and (b) R is aralkyl and R' is hydrogen or aralkyl, aryl, alkyl.

The term "acyl" refers to -C(O)R where R is alkyl and aryl. By way of example
25 only, acyl radicals include acetyl, pentanoyl, benzoyl, 4-hydroxybenzoyl, pivaloyl and 4-hydroxyphenylacetyl.

The term "carboxy esters" refers to -C(O)OR where R is alkyl, aryl, aralkyl, and alicyclic, all optionally substituted.

The term "carboxyl" refers to -C(O)OH.

30 The term "oxo" refers to =O attached to carbon or hetero atom.

The term "amino" refers to $-NRR'$ where R and R' are independently selected from hydrogen, alkyl, aryl, aralkyl and alicyclic, all except H are optionally substituted; and R and R' can form a cyclic ring system.

The term "halogen" or "halo" refers to -F, -Cl, -Br and -I.

5 The term "heterocyclic" and "heterocyclic alkyl" refer to cyclic groups containing at least one heteroatom. Suitable heteroatoms include oxygen, sulfur, and nitrogen. Heterocyclic groups may be attached through a nitrogen or through a carbon atom in the ring. Suitable heterocyclic groups include pyrrolidinyl, morpholino, morpholinoethyl, and pyridyl.

10 The term "phosphono" refers to $-PO_3R_2$, where R is selected from the group consisting of -H, alkyl, aryl, aralkyl, and alicyclic.

The term "sulfonyl" refers to $-SO_3R$, where R is H, alkyl, aryl, aralkyl, and alicyclic.

15 The term "acyloxy" refers to the ester group $-O-C(O)R$, where R is H, alkyl, alkenyl, alkynyl, aryl, aralkyl, or alicyclic.

The term "hydroxy" refers to the OH group

The term "thiol" refers to the SH group.

The term "alkylamino" refers to an amine- NR_2 wherein at least one R is an alkyl group.

20 The term " (C_1-C_8) alkylamino" refers to an alkylamino group having a single alkyl group, wherein the alkyl group has up to and including 8 carbon atoms. Suitable alkyl groups may be straight chain, branched, or cyclic and may be optionally substituted.

25 The term "dialkylamino" refers to an amine- NR_2 wherein both R groups are alkyl groups. Suitable alkyl groups may be straight chain, branched, or cyclic and may be optionally substituted.

The term " $di(C_1-C_8)$ alkylamino" refers to an amine- NR_2 wherein both R groups are alkyl groups. Suitable alkyl groups may be straight chain, branched, or cyclic and may be optionally substituted. Further, the two alkyl groups may be joined to form a ring comprising the nitrogen as a hetero atom.

30 The term "aminoalkyl-" refers to the group NR_2 -alk- wherein "alk" is an alkylene group and R is selected from H, alkyl, aryl, aralkyl, and alicyclic.

The term "alkylaminoalkyl-" refers to the group alkyl-NR-alk- wherein "alk" is an alkylene group and R is H or lower alkyl. "Lower alkylaminoalkyl-" refers to groups where each alkyl group is lower alkyl.

5 The term "arylaminoalkyl-" refers to the group aryl-NR-alk- wherein "alk" is an alkylene and R is H, alkyl, aryl, aralkyl, and alicyclic. In "lower arylaminoalkyl-", the alkyl group is lower alkyl.

The term "alkylaminoaryl-" refers to the group alkyl-NR-aryl- wherein "aryl" is a divalent group and R is H, alkyl, aralkyl, and alicyclic. In "lower alkylaminoaryl-", the alkyl group is lower alkyl.

10 The term "alkyloxyaryl-" refers to the group alkyl-O-aryl- wherein an "aryl" is a divalent group. In "lower alkyloxyaryl-", the alkyl group is lower alkyl.

The term "aryloxyalkyl-" refers to an alkylene group substituted with an aryloxy group.

15 The term "alkoxy" refers to an alkyl group as described above which also bears an oxygen substituent which is capable of covalent attachment to another hydrocarbon radical (such as, for example, methoxy, ethoxy and t-butoxy).

The term (C₁-C₈)alkoxy as used herein refers to an alkoxy group, wherein alkyl is a (C₁-C₈)alkyl.

20 The term "alkenyloxy" as used herein refers to the group -O-alkenyl wherein "alkenyl" is an alkenyl group.

The term (C₁-C₈)alkenyloxy as used herein refers to alkenyloxy, wherein the alkenyl is a (C₁-C₈)alkenyl.

The term "aryloxy" as used herein refers to the group Ar-O- wherein "Ar" is an aryl group.

25 The terms "alkylthio-" as used herein refers to the group alkyl-S-.

The term "(C₁-C₈)alkylthio" as used herein refers to the group alkyl-S- wherein alkyl has up to and including 8 carbon atoms.

30 The terms "amido" or "carboxamido" as used herein refers to NR₂-C(O)- and RC(O)-NR¹-, where R and R¹ include H, alkyl, aryl, aralkyl, and alicyclic. The term does not include urea, -NR-C(O)-NR-.

The term "aminocarboxamidoalkyl-" as used herein refers to the group NR₂-C(O)-N(R)-alk- wherein R includes H, alkyl, aryl, aralkyl, and alicyclic, and "alk" is an alkylene

group. "Lower aminocarboxamidoalkyl-" refers to such groups wherein "alk" is lower alkylene.

The term "heteroarylalkyl" refers to an alkyl group substituted with a heteroaryl group.

5 The term "perhalo" as used herein refers to groups wherein every C-H bond has been replaced with a C-halo bond. Suitable perhaloalkyl groups include -CF₃ and -CFC1₂. Suitable perhaloalkylene groups include -CF₂ and mixed halogen species, e.g., -CFC1.

The term "cyano" or "nitrile" as used herein refers to a -C≡N group.

The term "nitro" as used herein refers to a -NO₂ group.

10 The term "alkylsulfonyl" as used herein refers alkOS(O)₂- wherein alk is an alkyl.

The term "phosphonate" as used herein refers to the moiety P(=O)(OR)₂ wherein P(=O) designates a phosphorous oxo group, an wherein R may be an aliphatic group.

The term "ester" as used herein refers to a chemical moiety with formula -(R)_n-COOR', where R and R' are independently selected from the group consisting of alkyl or aryl and n is 0 or 1.

15 The term "amide" as used herein refers to a chemical moiety with formula -(R)_n-CONHR', where R and R' are independently selected from the group consisting of alkyl or aryl and n is 0 or 1.

20 The term "imide" as used herein refers to a chemical moiety with formula -(R)_n-CONR₂', where R and R' are independently selected from the group consisting of alkyl or aryl and n is 0 or 1.

The term "cross-coupling" as used herein refers a reaction between an electrophile and a nucleophile resulting in the formation of a new covalent carbon-carbon bond.

25 The term "organostannane" or "organostannane compound" as used herein refers to a compound comprising at least one Sn-R chemical bond, wherein R is an organic moiety. Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety. Preferred organostannanes are any which can engage in cross-coupling reactions.

The term "organozinc" or "organozinc compound" as used herein refers to a compound having at least one Zn-R chemical bond, wherein R is an organic moiety.

30 Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety.

Preferred organozinc compounds according to the present invention are those which engage in cross-coupling reactions:

The term "organoboron" or "organoboron compound" as used herein refers to a compound comprising at least one B-R chemical bond, wherein R is an organic moiety.

5 Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety. Preferred organoborons are those which engage in cross-coupling reactions.

The term "organoaluminum" or "organoaluminum compound" as used herein refers to a compound comprising at least one Al-R chemical bond, wherein R is an organic moiety. Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety. Preferred organoaluminums are those which engage in cross-coupling reactions.

10 The term "organomagnesium", "organomagnesium compound", "organomagnesium reagent" or "Grignard reagent" as used herein refers to a compound comprising at least one Mg-R chemical bond, wherein R is an organic moiety. Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety. Preferred organomagnesium compounds are those which engage in cross-coupling reactions

The term "organolithium" or "organolithium compound" as used herein refers to a compound comprising at least one Li-R chemical bond, wherein R is an organic moiety. Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety.

20 The term "organosilicon" or "organosilicon reagent" as used herein refers to a compound comprising at least one Si-R chemical bond, wherein R is an organic moiety. Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety.

The term "organocopper reagent" also known as "organocuprates" as used herein refers to a compound comprising at least one Cu-R chemical bond, wherein R is an organic moiety. Preferably R can be either an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety.

25 The term dialkylphosphite, refers to compound having a formula: $R_2P(=O)OR'$ wherein R and R' comprise aliphatic or aryl groups. Preferably R and R' are independently selected from the group consisting of an alkyl, alkenyl, aryl, heteroaryl, allyl or alkynyl moiety.

30 The term "aryl boronic acid" as used herein refers to compounds having a formula: $ArB(OH)_2$, wherein Ar refers to an aryl.

The term "metal halide" as used herein refers to any covalently or ionically bonded compound comprising at least one electropositive element and at least one electro-negative element. According to one aspect of the present invention, a metal halide is a source of a halide anion.

5 The term "addition of a transition metal catalyst" refers to the use of a transition metal catalyst to promote a desired chemical transformation. A transition metal catalyst comprises at least one transition metal and associated ligands. A non-catalytic or poorly catalytic transition metal can often be converted to a transition metal catalyst by the addition of ligands into the reaction mixture; such ligands include, by way of example
10 only, trialkylphosphines, triarylphosphines, and bis(diarylphosphino)L compounds, wherein L can be an alkylene, divalent aryl, divalent alkenyl, divalent alkynyl, divalent heteroalkyl, divalent heteroaryl, or a derivative of ferrocene. In one set of embodiments, ligands include 1,3-bis(diphenylphosphino)propane or 1,1'-bis(diphenylphosphanyl)ferrocene. Any particular transition metal catalyst may have
15 multiple ligands, not all of which need to be identical. In one embodiment, the transition metal is palladium. The new catalytically active species are also known as transition metal complexes. The skilled artisan will understand that the examples provided herein for transition metal catalyst, transition metal and ligand are for illustrative purposes only and that the scope of the present invention is not limited by the identity of the transition metal
20 catalyst, the transition metal, and the ligands. The catalyst and the optional ligand which together form a catalytically active species may be used at levels close to 100% relative to the reacting species, preferably <25% relative to the reacting species, more preferably <10% relative to the reacting species.

25 The term "covalently attached" as used herein refers to the presence of a covalent bond between two or more moieties.


 The term "nucleophile" as used herein is a compound or moiety that is reactive towards an electrophile so as to form a covalent between the nucleophile and electrophile. The terms "nucleophile" and "electrophile" have their usual meanings familiar to synthetic and/or physical organic chemistry. Carbon electrophiles typically comprise one or more
30 alkyl, alkenyl, alkynyl or aromatic carbon atom substituted with any atom or group having a Pauling electronegativity greater than that of hydrogen. Examples of preferred carbon electrophiles include but are not limited to carbonyls (especially aldehydes and ketones),

oximes, hydrazones, epoxides, aziridines, alkyl-, alkenyl-, and aryl halides, acyls, sulfonates, and perhalosulfonates. Other examples of carbon electrophiles include unsaturated carbons electronically conjugated with electron-withdrawing groups, examples being the β -carbon in α,β -unsaturated ketones or carbon atoms in fluorine substituted aryl groups. The skilled artisan will realize that the scope of the invention is not limited by the identity of the nucleophile.

The term "activated complex" as used herein refers to a moiety that is covalently bound to a surface-bound activator and which, as a result, has an increased chemical reactivity relative to its parent compound. According to one embodiment of the present invention an aryl alcohol compound is converted into a more reactive, surface-bound aryloxy moiety, i.e., an activated complex, upon reaction with a surface-bound activator of the present invention.

The term "hydride" or "hydride reagent" as used herein, refers to proton that can act as a nucleophile. Examples of hydride reagents familiar to the skilled artisan include but are not limited to NaBH_4 , LiAlH_4 , as well as any suitable transition-metal hydride. The hydride may also be formed in situ, or during the course of the reaction, by an interaction between one or more reagents. By way of example only, a transition metal-hydride is formed in situ in the presence of $\text{Pd}(\text{OAc})_2$, ligand and formic acid, as described in Example 3 below.

The term "enolizable ketone" refers to any ketone bearing a hydrogen on an "alpha carbon" (the carbon atom once removed from that bearing the oxo group), wherein said alpha proton may be readily removed by exogenous base or via intra molecular tautomerization:



The term "compound" as used herein refers to any identifiable molecule.

The term "conditions" refers to factors that can affect the outcome of a particular reaction and which can be controlled by the operator performing the reaction or sequence of reactions. Examples of such conditions include but are not limited to the length of time
5 that a set of reagents is allowed to interact with a substrate, the temperature, the solvent, the rates of addition of particular reagents, and the like. The skilled artisan will recognize that each particular set of reagents may have its own optimal set of "conditions".

The term "conditions sufficient" as used herein refers to those conditions that are adequate to produce the results desired.

10 The term "contacting" as used herein refers to any process whereby any reagent or combinations thereof is mixed, stirred, added to, shaken, dissolved, passed-over, passed through, another said reagent under conditions such that two or more reagents can undergo a chemical reaction or transformation.

The term "reagent" as used herein refers to any chemical compound used alone or
15 in combination with a different chemical compound to produce a desired chemical reaction. The term "reagent" includes all catalysts (transition metal based or otherwise), ligands, acids, bases and other materials that are added to a reaction mixture in order to provide the desired result.

The term "moiety" as used herein refers to a specific portion of a molecule, usually
20 complex, that has a characteristic chemical or property or reactivity.

The term "linking group" as used herein refers to the entire chain of atoms linking an activator portion with a solid or semi-solid support.

General

25 The present invention provides a variety of reagents, particularly support-bound activators, that have utility in the area of combinatorial synthesis. Each reagent includes an activator portion that serves as a reactive center and a linking group component that serves to provide a robust linkage between the support and the activator portion. The linking group components further include an activator enhancing portion that serves to
30 increase the reactivity of the activator portion and suitable spacer that provides sufficient distance between the activator portion and the support. The present invention also provides

for activated supports on which construction of a target or library of targets takes place; the targets can also be cleaved from the activated support to liberate the desired compound from the activated support.

One important aspect of the present invention is the use of the activated supports described herein as traceless linkers, as that term is understood by those skilled in the art. As described above, the term "traceless linker" has been used to describe a strategy of releasing compounds from a solid support with little or no trace of the original point of attachment. See James, *Tetrahedron Lett.*, 1999, 55, 4855; Andres, *et al.*, *Curr. Opin. Chem. Biol.*, 1998, 2, 353; Reitz, *Curr. Opin. Drug Discovery Dev.*, 1999, 2, 358; and Zaragoza, *Angew. Chem., Int. Ed.* 2000, 39, 2077. As demonstrated in Examples 3, 4 and 5, and in Tables 1, 2, and 3, the activated supports of the present invention will have wide utility as traceless linkers in solid phase organic chemistry. As one skilled in the art will realize, these examples are merely illustrative and the activated supports of the present invention are fully expected to act as traceless linkers in any reaction that can cleave a triflate-carbon bond.

In addition, any of the reactions described herein, as well as any reaction known in the art of organic chemistry that can retain the integrity of a triflate-carbon bond, can be used to construct either a target or a library of targets on the activated supports of the present invention. Accordingly a preferred aspect of the present invention is a support-activated target group. In addition, another preferred aspect of the present invention is a library of support-activated target groups.

The support-bound activators of the present invention also serve to activate certain centers toward such reactions as reductions, Suzuki couplings, Stille couplings, Heck couplings, Buchwald reactions, CO insertions, CN insertions, carbon-sulfur bond formation, and others.

The use of triflates and nonaflates as precursors for aryl and vinyl cations has been widely recognized. See, for example, Ritter, *Synthesis*, 8:735-762 (1993). Briefly, an oxygen atom on an aryl or vinyl group (e.g., as a phenol or an enolizable ketone) can be activated as a triflate ester (trifluoromethane sulfonate ester) or a related nonaflate ester towards a subsequent reduction or cross-coupling that gives rise to a variety of substituted aromatic compounds or olefinic compounds.

It would be desirable to conduct the perfluorosulfonyl-directed transformations on solid phase and take advantage of the versatile synthetic transformations known for vinyl and aryl triflates (see, Ritter, *ibid.*). For example, the reductive cleavage of a polymer-supported aryl triflate could lead to deoxygenation of phenols without leaving a trace of the phenolic hydroxy group as a point of attachment to the polymer, or resin. Wustrow and coworkers (*Tetrahedron Lett.* 39:3651 (1998)) reported the reductive cleavage of electron poor arylsulfonates from an ion-exchange resin based sulfonyl linker under stringent cleavage conditions (140°C for 12 h), but to date, there have been no reports of polymer-supported triflates and nonaflates that could be used for activating hydroxy groups toward a variety of synthetic transformations.

NAFION™ resin is well known, yet suffers from having poor swelling properties and is difficult to activate. The inability of NAFION™ to swell in the presence of common organic solvents prevents the vast majority of surface-bound reactive groups from reacting with compounds in solution. As a result, NAFION™ has not found much utility in solid-phase organic synthesis or in other uses that require more than a catalytic amount of available surface-bound reactive groups.


What is needed in the art are support-bound activators that have improved swelling properties and that are able to participate in perfluorosulfonate-based activation/displacement chemistry. Such support-bound activators and the related activated supports are provided below.

The present invention also provides for strongly acidic supports which find application as catalysts or scavenger resins, particularly those having increased swelling properties. In addition, the present invention provides for silylating supports which find application in analytical chemistry or in other areas in which a silylated compound is desired.

Description of the Embodiments

Support-Bound Activators

In one aspect, the present invention provides a support-bound activator having the formula:



In formula I, the letter L represents a linking group component; the letter X represents F, Cl, trisubstituted silyloxy or OH. The support-bound activator is attached to a solid or semi-solid support by at least one covalent bond.

Turning first to the solid or semi-solid support, the present invention is useful in a variety of solid-phase synthesis applications and, accordingly, a variety of supports find utility in this aspect of the invention. Typical solid supports include, but are not limited to, cross-linked divinylbenzene-styrene (polystyrene), controlled pore glass (CPG), polyacrylamides, poly(ethyleneglycol)monomethyl ether and poly(ethylene glycol) (PEG), silica gel, cellulose, acrylic acid grafted polypropylene, and the like. Additionally, the solid support contains a reactive moiety suitable for attaching the linking group component. Suitably reactive moieties include, for example, a carboxylic acid, alcohol, amine, halomethyl and the like which is used to covalently attach the linking group component during construction of the present support-bound activators. Many of these supports are available as functional polymers having reactive groups. Examples of such supports, include, by way of example, Acryloyl Wang resin, REM resin, Vinyl polystyrene, Vinylsulfonylmethyl polystyrene, (3-Formylindolyl)acetamidomethyl polystyrene, 2-(3,5-Dimethoxy-4-formylphenoxy)ethoxymethyl polystyrene, 2-(4-Formyl-3-methoxyphenoxy)ethyl polystyrene, 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin, 4-Benzoyloxybenzaldehyde polystyrene, Aldehyde Wang resin, Formylpolystyrene, 1% DVB, NovaSyn® TG acetal resin, Polystyrene-CHO, Carboxypolystyrene, NovaSyn® TG carboxy resin, Polystyrene-COOH, 4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxy resin, 4-Methylbenzhydramine resin HCl, 4-Methylbenzhydramine resin HCl, 9-Fmoc-amino-xanthen-3-yloxy, 9-Fmoc-amino-xanthen-3-yloxy, Amino-(4-

methoxyphenyl)methyl polystyrene, Ethylamino-xanthen-3-yloxy-Merrifield resin, NovaSyn® TG Sieber resin, NovaSyn® TGR resin, Rink Amide AM resin, Rink Amide MBHA resin, Rink amide NovaGel™, Rink amide PEGA resin, Rink Amide resin, Sieber Amide resin, Sieber Ethylamide resin, Amino methyl resin, Amino PEGA resin,

5 Aminomethyl NovaGel™, Aminomethylated polystyrene, N-Methylaminomethyl polystyrene, 4-Fmoc-hydrazinobenzoyl AM resin, 1H-Benzotriazole polystyrene, Benzotriazole-5-carbamidomethyl polystyrene, N-Fmoc-N-methoxy-β-alanine AM resin, Weinreb AM resin, 4-Sulfamylbenzoyl AM resin, (±)-1-(2,3-Isopropylidene)glycerol polystyrene, (±)-2,2-Dimethyldioxolan-4-methoxymethyl polystyrene, (±)-1-Glycerol

10 polystyrene, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid AM resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid BHA resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid MBHA resin, 4-Hydroxymethylphenoxyacetyl NovaGel™,

15 4-Hydroxymethylphenoxyacetyl PEGA resin, HMP resin, HMPA-NovaGel™, HMPB-AM resin, HMPB-BHA resin, HMPB-MBHA resin, Hydroxy-(2-chlorophenyl)methyl polystyrene, Hydroxymethylpolystyrene, NovaSyn® TG HMP resin, p-Benzyloxybenzyl Alcohol resin, Polystyrene-CH₂OH, Rink Acid resin, Trichloroacetimidate Wang resin, Wang resin, 4-Hydroxymethylbenzoic acid AM resin, 4-Hydroxymethylbenzoic acid

20 NovaGel™, 4-Hydroxymethylbenzoic acid PEGA resin, 4-Hydroxyphenylsulfanylmethyl polystyrene, 9-(Hydroxymethyl)fluorene-4-carboxamidomethyl polystyrene, HESM polystyrene, HMBA-AM resin, HMBA-NovaGel™, HMBA-PEGA resin, Hydroxyethylsulfanylmethyl polystyrene, NovaSyn® TG HMBA resin, NovaSyn® TG hydroxy resin, Oxime resin, Aminoethyl photolinker resin, Hydroxyethyl photolinker

25 resins, Hydroxymethyl photolinker resins, 3-[4-(Tritylmercapto)phenyl]propionyl AM resin, Mercaptomethyl polystyrene, NovaSyn® TG tritylthiol resin, Thiol 2-chlorotrityl resin, Thiol 4-methoxytrityl resin, (4-Bromophenyl)diisopropylsilyloxymethyl polystyrene, (4-Formylphenyl)diisopropylsilyloxymethyl polystyrene, (4-Trityloxylphenyl)diisopropylsilyloxymethyl polystyrene. Solid supports also include

30 TENTAGEL™, HYPOGEL™, JANDAJEL™, AND ARGOGEL™. Other solid supports include PEGylated polystyrene (polystyrene derivatized with polyethylene glycol), Tentagel-NH₂ resin, and derivatized Tentagel-NH₂ resin (e.g., by treatment with acetyl

chloride followed by reduction with LiAlH_4 to provide Tentagel-NHEt resin). See also the Novabiochem Catalogue 2000 for additional resins and immobilized functional groups.



Such supports may take any size, shape or form, including particulate and non-particulate forms or shapes, spheres, disks, pellets, sheets, plugs, pins, crowns, lanterns, in
5 beaded and non-beaded forms, resins, gels, microspheres, as well as amorphous forms and shapes. Embodiments of particulate supports, include beads, pellets, disks, amorphous particles, or other conventional forms. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-through devices. In a one embodiment, the diameter of the
10 particulate support is 20-2000 micron, preferably 75-500 micron, more preferably 100-200 micron. As one skilled in the art will readily recognize, the scope of the present invention is not limited to the size, form, or shape of the solid or semi-solid support.

The linking group component can have a variety of structures. The linking group component is one which provides suitable spacing for the activator portion ($-\text{CF}_2-\text{SO}_2-\text{X}$)
15 to interact freely with molecules or reactive components exposed to the activator portion. The linking group component is preferably 6-50 atoms long, more preferably 8-40 atoms long, even more preferably 8-30 atoms long, and yet more preferably 8-20 atoms long, thus providing sufficient exposure for the attached activator portion. Additionally, the linking group component, prior to attachment to the support, will have a attaching portion
20 and a longer chain portion. The attaching portion is that part of the linking group component which can be directly attached to the solid support. This portion can be attached to the solid support via carbon-carbon bonds using, for example, supports having exposed (poly)trifluorochloroethylene moieties, or preferably, by siloxane bonds (using, for example, glass or silicon oxide as the solid support). Siloxane bonds the support are
25 formed in one embodiment via reactions of attaching portions bearing trichlorosilyl or trialkoxysilyl groups. The attaching groups will also have a site for attachment of the longer chain portion. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl.

One skilled in the art will recognize that many additional methods of attaching
30 linkers to solid and semi-solid supports exist. One method uses an amino resin to which an acid-bearing linker is attached via conventional techniques. Another method is the use of aryl ether linkages by coupling a phenol to a Merrifield (or equivalent) resin.

The longer chain portion can be any of a variety of molecules which are inert to the subsequent conditions used in the activator reactions described in further detail below. These longer chain portions can be ethylene glycol oligomers containing 2-14 monomer units, or more preferably 2-10 monomer units, and even more preferably 2-8 monomer units; in addition, the longer chain portions can be diamines, diacids, amino acids, peptides, or combinations thereof. In some embodiments, the longer chain portion also comprises an activator enhancing portion, i.e., a portion that increases the reactivity of the activator relative to an alkylene or ethylene glycol linking group. More particularly, an activator enhancing portion is one that provides additional electron withdrawing character to the activator portion (e.g., the $-\text{CF}_2-\text{SO}_2-\text{X}$ portion).

In one group of embodiments, X is a member selected from the group consisting of F and Cl. In another group of embodiments, L comprises an activator enhancing portion selected from the group consisting of:



wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, and NR^1 ; Z is a member selected from the group consisting of a chemical bond or CO; each R^2 is independently a member selected from the group consisting of hydrogen, halogen, $(\text{C}_1\text{--C}_8)$ alkyl, $(\text{C}_1\text{--C}_8)$ alkoxy, $(\text{C}_2\text{--C}_8)$ heteroalkyl, $(\text{C}_1\text{--C}_8)$ alkylthio, $(\text{C}_1\text{--C}_8)$ alkylamino, $\text{di}(\text{C}_1\text{--C}_8)$ alkylamino, cyano, nitro and $(\text{C}_1\text{--C}_8)$ alkylsulfonyl; the subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or

can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).


In another embodiment, the support-bound activator is available in kit form for use in solid phase organic chemistry, as a reagent or catalyst in solution phase organic chemistry, as a scavenger resin in solution phase organic chemistry, as a silylating agent for use in analytical chemistry, and in particular, in chromatography, and as a reagent for the production of PET-ready molecules.

In one group of of embodiments, the support-bound activators of the present invention are:




wherein the symbol R^1 and the subscripts g, i, j, k and m all have the meanings provided above.

In another group of embodiments, the support-bound activators have a formula selected from:



and



In each of formulae A and B, X can be F, Cl, trisubstituted silyloxy, or OH; Q is O; Z is a chemical bond or C=O; Y is O-support or NR_1 -support wherein R_1 is H, $(\text{C}_1\text{--C}_8)$ alkyl or aryl and the support is a PEG-modified polystyrene or a Merrifield resin; and each R^2 is as defined more generally above. The subscripts in formulae A and B are as follows: 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11; 'h' is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8; 'i' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, and 8; 'j' is an integer selected from the group consisting of 1, 2, 3, and 4; 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; 'm' is an integer selected from the group consisting

of 2 and 3; 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4; and 'q' is an integer selected from the group consisting of 1, and 2.


In one particular embodiment in formulae A and B, X is F; Q is O; Z is C=O, Y is NH-support wherein the support is a PEG-modified polystyrene; and each R² is H.

5

Activated Supports

In another aspect, the present invention provides an activated support comprising a solid or semi-solid support; and at least one support-bound activator having the formula:

10



wherein L is a linking group component; X is a member selected from the group consisting of F, Cl, OH, and trisubstituted silyloxy; and wherein the support-bound activator is covalently attached to the solid or semi-solid support.

Turning first to the solid or semi-solid support, the present invention is useful in a variety of solid-phase synthesis applications and, accordingly, a variety of supports find utility in this aspect of the invention. Typical solid supports include, but are not limited to, cross-linked divinylbenzene-styrene (polystyrene), controlled pore glass (CPG), polyacrylamides, poly(ethyleneglycol)monomethyl ether and poly(ethylene glycol) (PEG), silica gel, cellulose, acrylic acid grafted polypropylene, and the like. Additionally, the solid support contains a reactive moiety suitable for attaching the linking group component. Suitably reactive moieties include, for example, a carboxylic acid, alcohol, amine, halomethyl and the like which is used to covalently attach the linking group component during construction of the present support-bound activators. Many of these supports are available as functional polymers having reactive groups. Examples of such supports, include, by way of example, Acryloyl Wang resin, REM resin, Vinyl polystyrene, Vinylsulfonylmethyl polystyrene, (3-Formylindolyl)acetamidomethyl polystyrene, 2-(3,5-Dimethoxy-4-formylphenoxy)ethoxymethyl polystyrene, 2-(4-Formyl-3-methoxyphenoxy)ethyl polystyrene, 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin, 4-Benzoyloxybenzaldehyde polystyrene, Aldehyde Wang resin, Formylpolystyrene, 1% DVB, NovaSyn® TG acetal resin, Polystyrene-CHO, Carboxypolystyrene, NovaSyn® TG

carboxy resin, Polystyrene-COOH, 4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxymethyl resin, 4-Methylbenzhydrylamine resin HCl, 4-Methylbenzhydrylamine resin HCl, 9-Fmoc-amino-xanthen-3-yloxy, 9-Fmoc-amino-xanthen-3-yloxy, Amino-(4-methoxyphenyl)methyl polystyrene, Ethylamino-xanthen-3-yloxy-Merrifield resin,

5 NovaSyn® TG Sieber resin, NovaSyn® TGR resin, Rink Amide AM resin, Rink Amide MBHA resin, Rink amide NovaGel™, Rink amide PEGA resin, Rink Amide resin, Sieber Amide resin, Sieber, Ethylamide resin, Amino methyl resin, Amino PEGA resin, Aminomethyl NovaGel™, Aminomethylated polystyrene, N-Methylaminomethyl polystyrene, 4-Fmoc-hydrazinobenzoyl AM resin, 1H-Benzotriazole polystyrene,

10 Benzotriazole-5-carbamidomethyl polystyrene, N-Fmoc- N-methoxy-β-alanine AM resin, Weinreb AM resin, 4-Sulfamylbenzoyl AM resin, (±)-1-(2,3-Isopropylidene)glycerol polystyrene, (±)-2,2-Dimethyldioxolan-4-methoxymethyl polystyrene, (±)-1-Glycerol polystyrene, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxymethyl resin, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxymethyl resin, 4-(2',4'-Dimethoxyphenyl-

15 hydroxymethyl)-phenoxymethyl resin, 4-Hydroxymethyl-3-methoxyphenoxymethylbutyric acid AM resin, 4-Hydroxymethyl-3-methoxyphenoxymethylbutyric acid BHA resin, 4-Hydroxymethyl-3-methoxyphenoxymethylbutyric acid MBHA resin, 4-Hydroxymethylphenoxymethylbutyric acid NovaGel™, 4-Hydroxymethylphenoxymethylbutyric acid PEGA resin, HMP resin, HMPA-NovaGel™, HMPB-AM resin, HMPB-BHA resin, HMPB-MBHA resin, Hydroxy-(2-chlorophenyl)methyl

20 polystyrene, Hydroxymethylpolystyrene, NovaSyn® TG HMP resin, p-Benzoyloxymethyl Alcohol resin, Polystyrene-CH₂OH, Rink Acid resin, Trichloroacetimidate Wang resin, Wang resin, 4-Hydroxymethylbenzoic acid AM resin, 4-Hydroxymethylbenzoic acid NovaGel™, 4-Hydroxymethylbenzoic acid PEGA resin, 4-Hydroxyphenylsulfanylmethyl polystyrene, 9-(Hydroxymethyl)fluorene-4-carboxamidomethyl polystyrene, HESM

25 polystyrene, HMBA-AM resin, HMBA-NovaGel™, HMBA-PEGA resin, Hydroxyethylsulfanylmethyl polystyrene, NovaSyn® TG HMBA resin, NovaSyn® TG hydroxy resin, Oxime resin, Aminoethyl photolinker resin, Hydroxyethyl photolinker resins, Hydroxymethyl photolinker resins, 3-[4-(Tritylmercapto)phenyl]propionyl AM resin, Mercaptomethyl polystyrene, NovaSyn® TG tritylthiol resin, Thiol 2-chlorotrityl

30 resin, Thiol 4-methoxytrityl resin, (4-Bromophenyl)diisopropylsilyloxymethyl polystyrene, (4-Formylphenyl)diisopropylsilyloxymethyl polystyrene, (4-Trityloxymethylphenyl)diisopropylsilyloxymethyl polystyrene. Solid supports also include

TENTAGEL™, HYPOGEL™, JANDAJEL™, AND ARGOGEL™. Other solid supports include PEGylated polystyrene (polystyrene derivatized with polyethylene glycol), Tentagel-NH₂ resin, and derivatized Tentagel-NH₂ resin (e.g., by treatment with acetyl chloride followed by reduction with LiAlH₄ to provide Tentagel-NHEt resin). See also the
5 Novabiochem Catalogue 2000 for additional resins and immobilized functional groups.


Such supports may take any size, shape or form, including particulate and non-particulate forms or shapes, spheres, disks, pellets, sheets, plugs, pins, crowns, lanterns, in beaded and non-beaded forms, resins, gels, microspheres, as well as amorphous forms and shapes. Embodiments of particulate supports, include beads, pellets, disks, amorphous
10 particles, or other conventional forms. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-through devices. In a one embodiment, the diameter of the particulate support is 20-2000 micron, preferably 75-500 micron, more preferably 100-200 micron. As one skilled in the art will readily recognize, the scope of the present invention
15 is not limited to the size, form, or shape of the solid or semi-solid support.

The linking group component can have a variety of structures. The linking group component is one which provides suitable spacing for the activator portion (-CF₂-SO₂-X) to interact freely with molecules or reactive components exposed to the activator portion. The linking group component is preferably 6-50 atoms long, more preferably 8-40 atoms
20 long, even more preferably 8-30 atoms long, and yet more preferably 8-20 atoms long, thus providing sufficient exposure for the attached activator portion. Additionally, the linking group component, prior to attachment to the support, will have a attaching portion and a longer chain portion. The attaching portion is that part of the linking group component which can be directly attached to the solid support. This portion can be
25 attached to the solid support via carbon-carbon bonds using, for example, supports having exposed (poly)trifluorochloroethylene moieties, or preferably, by siloxane bonds (using, for example, glass or silicon oxide as the solid support). Siloxane bonds the support are formed in one embodiment via reactions of attaching portions bearing trichlorosilyl or trialkoxysilyl groups. The attaching groups will also have a site for attachment of the
30 longer chain portion. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl.

One skilled in the art will recognize that many additional methods of attaching linkers to solid and semi-solid supports exist. One method uses an amino resin to which an acid-bearing linker is attached via conventional techniques. Another method is the use of aryl ether linkages by coupling a phenol to a Merrifield (or equivalent) resin.

5 The longer chain portion can be any of a variety of molecules which are inert to the subsequent conditions used in the activator reactions described in further detail below. These longer chain portions can be ethylene glycol oligomers containing 2-14 monomer units, or more preferably 2-10 monomer units, and even more preferably 2-8 monomer units; in addition, the longer chain portions can be diamines, diacids, amino acids, peptides, or combinations thereof. In some embodiments, the longer chain portion also
10 comprises an activator enhancing portion, i.e., a portion that increases the reactivity of the activator relative to an alkylene or ethylene glycol linking group. More particularly, an activator enhancing portion is one that provides additional electron withdrawing character to the activator portion (e.g., the $-\text{CF}_2-\text{SO}_2-\text{X}$ portion).

15 In one group of embodiments, X is a member selected from the group consisting of F and Cl. In another group of embodiments, L comprises an activator enhancing portion selected from the group consisting of:



20 wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, and NR^1 ; Z is a member selected from the group consisting of a chemical bond or CO; each R^2 is independently a member selected from the group consisting of hydrogen, halogen, $(\text{C}_1\text{-C}_8)\text{alkyl}$, $(\text{C}_1\text{-C}_8)\text{alkoxy}$, $(\text{C}_2\text{-C}_8)\text{heteroalkyl}$, $(\text{C}_1\text{-C}_8)\text{alkylthio}$, $(\text{C}_1\text{-C}_8)\text{alkylamino}$, $\text{di}(\text{C}_1\text{-C}_8)\text{alkylamino}$, cyano, nitro and $(\text{C}_1\text{-C}_8)\text{alkylsulfonyl}$; the subscript
25 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9,


and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or
 5 can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).

In one group of of embodiments, the support-bound activators of the present invention are:




10 wherein the symbol R^1 and the subscripts g, i, j, k and m all have the meanings provided above.

In another group of embodiments, the support-bound activators have a formula selected from:



and



15 In each of formulae A and B, X can be F, Cl, trisubstituted silyloxy, or OH; Q is O; Z is a chemical bond or C=O; Y is O-support or NR_1 -support wherein R_1 is H, $(\text{C}_1\text{--C}_8)$ alkyl or aryl and the support is a PEG-modified polystyrene or a Merrifield resin; and each R^2 is as defined more generally above. The subscripts in formulae A and B are as follows: 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9,
 20 10, and 11; 'h' is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8; 'i' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, and 8; 'j' is an integer selected from the group consisting of 1, 2, 3, and 4; 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; 'm' is an integer selected from the group consisting of 2 and 3; 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4; and 'q'
 25 is an integer selected from the group consisting of 1, and 2.


In one particular embodiment in formulae A and B, X is F; Q is O; Z is C=O, Y is NH-support wherein the support is a PEG-modified polystyrene; and each R² is H.

In still other embodiments, the activated support comprises a plurality of support-bound activators, with a concentration of at least 1 nmol support-bound activators per gram of activated support, more preferably, at least 1 μ mol support-bound activators per gram of activated support, and even more preferably, at least 1 mmol support-bound activators per gram of activated support.

In another embodiment, the activated support is available in kit form for use in solid phase organic chemistry, as a reagent or catalyst in solution phase organic chemistry, as a scavenger resin in solution phase organic chemistry, as a silylating agent for use in analytical chemistry, and in particular, in chromatography, and as a reagent for the production of PET-ready molecules.

Support-Activated Targets

In another aspect, the present invention provides a support-activated target comprising a solid or semi-solid support; an activating group covalently attached to the solid or semi-solid support, wherein the activating group has the formula:



wherein L is an linking group component; and a target group covalently attached to the solid or semi-solid support; wherein the target group can be cleaved from the activating group by a nucleophile.

Turning first to the solid or semi-solid support, the present invention is useful in a variety of solid-phase synthesis applications and, accordingly, a variety of supports find utility in this aspect of the invention. Typical solid supports include, but are not limited to, cross-linked divinylbenzene-styrene (polystyrene), controlled pore glass (CPG), polyacrylamides, poly(ethyleneglycol)monomethyl ether and poly(ethylene glycol) (PEG), silica gel, cellulose, acrylic acid grafted polypropylene, and the like. Additionally, the solid support contains a reactive moiety suitable for attaching the linking group component. Suitably reactive moieties include, for example, a carboxylic acid, alcohol,

amine, halomethyl and the like which is used to covalently attach the linking group component during construction of the present support-bound activators. Many of these supports are available as functional polymers having reactive groups. Examples of such supports, include, by way of example, Acryloyl Wang resin, REM resin, Vinyl

5 polystyrene, Vinylsulfonylmethyl polystyrene, (3-Formylindolyl)acetamidomethyl polystyrene, 2-(3,5-Dimethoxy-4-formylphenoxy)ethoxymethyl polystyrene, 2-(4-Formyl-3-methoxyphenoxy)ethyl polystyrene, 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin, 4-Benzyloxybenzaldehyde polystyrene, Aldehyde Wang resin, Formylpolystyrene, 1% DVB, NovaSyn® TG acetal resin, Polystyrene-CHO, Carboxypolystyrene, NovaSyn® TG

10 carboxy resin, Polystyrene-COOH, 4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxy resin, 4-Methylbenzhydramine resin HCl, 4-Methylbenzhydramine resin HCl, 9-Fmoc-amino-xanthen-3-yloxy, 9-Fmoc-amino-xanthen-3-yloxy, Amino-(4-methoxyphenyl)methyl polystyrene, Ethylamino-xanthen-3-yloxy-Merrifield resin, NovaSyn® TG Sieber resin, NovaSyn® TGR resin, Rink Amide AM resin, Rink Amide

15 MBHA resin, Rink amide NovaGel™, Rink amide PEGA resin, Rink Amide resin, Sieber Amide resin, Sieber Ethylamide resin, Amino methyl resin, Amino PEGA resin, Aminomethyl NovaGel™, Aminomethylated polystyrene, N-Methylaminomethyl polystyrene, 4-Fmoc-hydrazinobenzoyl AM resin, 1H-Benzotriazole polystyrene, Benzotriazole-5-carbamidomethyl polystyrene, N-Fmoc- N-methoxy-β-alanine AM resin,

20 Weinreb AM resin, 4-Sulfamylbenzoyl AM resin, (±)-1-(2,3-Isopropylidene)glycerol polystyrene, (±)-2,2-Dimethyldioxolan-4-methoxymethyl polystyrene, (±)-1-Glycerol polystyrene, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid AM

25 resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid BHA resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid MBHA resin, 4-Hydroxymethylphenoxyacetyl NovaGel™, 4-Hydroxymethylphenoxyacetyl PEGA resin, HMP resin, HMPA-NovaGel™, HMPB-AM resin, HMPB-BHA resin, HMPB-MBHA resin, Hydroxy-(2-chlorophenyl)methyl polystyrene, Hydroxymethylpolystyrene, NovaSyn® TG HMP resin, p-Benzyloxybenzyl

30 Alcohol resin, Polystyrene-CH₂OH, Rink Acid resin, Trichloroacetimidate Wang resin, Wang resin, 4-Hydroxymethylbenzoic acid AM resin, 4-Hydroxymethylbenzoic acid NovaGel™, 4-Hydroxymethylbenzoic acid PEGA resin, 4-Hydroxyphenylsulfanylmethyl

polystyrene, 9-(Hydroxymethyl)fluorene-4-carboxamidomethyl polystyrene, HESM polystyrene, HMBA-AM resin, HMBA-NovaGel™, HMBA-PEGA resin, Hydroxyethylsulfanylmethyl polystyrene, NovaSyn® TG HMBA resin, NovaSyn® TG hydroxy resin, Oxime resin, Aminoethyl photolinker resin, Hydroxyethyl photolinker resins, Hydroxymethyl photolinker resins, 3-[4-(Tritylmercapto)phenyl]propionyl AM resin, Mercaptomethyl polystyrene, NovaSyn® TG tritylthiol resin, Thiol 2-chlorotrityl resin, Thiol 4-methoxytrityl resin, (4-Bromophenyl)diisopropylsilyloxymethyl polystyrene, (4-Formylphenyl)diisopropylsilyloxymethyl polystyrene, (4-Trityloxyphenyl)diisopropylsilyloxymethyl polystyrene. Solid supports also include TENTAGEL™, HYPOGEL™, JANDAJEL™, AND ARGOGEL™. Other solid supports include PEGylated polystyrene (polystyrene derivatized with polyethylene glycol), Tentagel-NH₂ resin, and derivatized Tentagel-NH₂ resin (e.g., by treatment with acetyl chloride followed by reduction with LiAlH₄ to provide Tentagel-NHEt resin). See also the Novabiochem Catalogue 2000 for additional resins and immobilized functional groups.

Such supports may take any size, shape or form, including particulate and non-particulate forms or shapes, spheres, disks, pellets, sheets, plugs, pins, crowns, lanterns, in beaded and non-beaded forms, resins, gels, microspheres, as well as amorphous forms and shapes. Embodiments of particulate supports, include beads, pellets, disks, amorphous particles, or other conventional forms. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-through devices. In a one embodiment, the diameter of the particulate support is 20-2000 micron, preferably 75-500 micron, more preferably 100-200 micron. As one skilled in the art will readily recognize, the scope of the present invention is not limited to the size, form, or shape of the solid or semi-solid support.



The linking group component can have a variety of structures. The linking group component is one which provides suitable spacing for the target group to interact freely with molecules or reactive components exposed to the target group. The linking group component is preferably 6-50 atoms long, more preferably 8-40 atoms long, even more preferably 8-30 atoms long, and yet more preferably 8-20 atoms long, thus providing sufficient exposure for the attached target group. Additionally, the linking group component, prior to attachment to the support, will have a attaching portion and a longer chain portion. The attaching portion is that part of the linking group component which can

be directly attached to the solid support. This portion can be attached to the solid support via carbon-carbon bonds using, for example, supports having exposed (poly)trifluorochloroethylene moieties, or preferably, by siloxane bonds (using, for example, glass or silicon oxide as the solid support). Siloxane bonds the support are
 5 formed in one embodiment via reactions of attaching portions bearing trichlorosilyl or trialkoxysilyl groups. The attaching groups will also have a site for attachment of the longer chain portion. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl.

One skilled in the art will recognize that many additional methods of attaching
 10 linkers to solid and semi-solid supports exist. One method uses an amino resin to which an acid-bearing linker is attached via conventional techniques. Another method is the use of aryl ether linkages by coupling a phenol to a Merrifield (or equivalent) resin.

The longer chain portion can be any of a variety of molecules which are inert to the subsequent conditions described in further detail below. These longer chain portions can
 15 be ethylene glycol oligomers containing 2-14 monomer units, or more preferably 2-10 monomer units, and even more preferably 2-8 monomer units; in addition, the longer chain portions can be diamines, diacids, amino acids, peptides, or combinations thereof. In some embodiments, the longer chain portion also comprises an activator enhancing portion, i.e., a portion that increases the reactivity of the target group relative to an
 20 alkylene or ethylene glycol linking group. More particularly, an activator enhancing portion is one that provides additional electron withdrawing character to the target group.

In another group of embodiments, L comprises an activator enhancing portion selected from the group consisting of:



25 wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, and NR^1 ; Z is a member selected from the group consisting of a chemical bond or CO;


each R² is independently a member selected from the group consisting of hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro and (C₁-C₈)alkylsulfonyl; the subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).

In still other embodiments, each solid or semi-solid support comprises a plurality of target groups, with a density of at least 1 nmol target groups per gram of solid or semi-solid support, more preferably, at least 1 μ mol target groups per gram of solid or semi-solid support, and even more preferably, at least 1 mmol target groups per gram of solid or semi-solid support.

In other embodiments, the support-activated target can be cleaved from the solid or semi-solid support by a reagent. In one group of embodiments, the reagent comprises a nucleophile; the cleavage step may also be promoted by a transition metal catalyst. Once cleaved from the support, the resulting compound can be isolated and characterized by methods standard to the synthesis of organic compounds. As one skilled in the art will readily recognize, the present invention is not limited by the particular use of the resulting compound.

Libraries of Support-Activated Targets

In another aspect, the present invention provides a library of support-activated targets comprising a plurality of support-activated target members, wherein each support-activated target member further comprises a solid or semi-solid support; an activating group covalently attached to the solid or semi-solid support, wherein the activating group has the formula:



wherein L is an linking group component; and a target group covalently attached to the activating group; wherein the target group of at least one support-activated target member in the library is different from the target group of at least one other support-activated target member in the library.

The description of the support-activated targets presented above is incorporated into this aspect of the present invention. As one skilled in the art will readily recognize, the present invention is not limited by the number of different support-activated target members in the library. Each support-activated target member of the library can be cleaved from its solid or semi-solid support by a reagent. In one set of embodiments the reagent comprises a nucleophile; the cleavage step may also be promoted by a transition metal catalyst. The support-activated targets and the resulting compounds can be used for a variety of purposes, including assays, screens, analysis, and testing. As one skilled in the art will readily recognize, the present invention is not limited by the particular use of the resulting compounds.

Linker Reagents



In view of the above, the present invention further provides linker reagents having the formula:



wherein L is a linking group component; and A is an attaching group.

In one embodiment, the attaching group is a member selected from the group consisting of NH_2 , NHR , CO_2H , CO_2R , $\text{C}(\text{O})\text{Cl}$, OH , SH , and protected forms thereof, wherein each R is a member independently selected from the group consisting of substituted $(\text{C}_1\text{-C}_8)\text{alkyl}$, unsubstituted $(\text{C}_1\text{-C}_8)\text{alkyl}$, substituted aryl, and unsubstituted aryl. In one embodiment, A is NH_2 or $\text{CH}_2\text{-halogen}$.

In further embodiments, L is selected from:




wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, NR¹; Z is a member selected from the group consisting of a chemical bond or CO; each R² is independently a member selected from the group consisting of hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro and (C₁-C₈)alkylsulfonyl; the subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).

In one group of embodiments, the linker reagent has the formula:



wherein the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; and the subscript k is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8. In one particular embodiment, the subscript j is 1 and the subscript k is 1.


In other embodiments, the linker reagent has the formula:



wherein the subscript g is an integer selected from the group consisting of 3, 4, 5, and 6; and the subscript i is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5,


5 6, 7, 8, 9, and 10.

In other embodiments, the linker reagent has the formula:




10

Other embodiments are those represented by formulae C and D:



and



in which X is F, Cl, trisubstituted silyloxy or OH; Q is O; A is C(O)Cl, CO₂H or OH; and each R² is as defined above. The subscripts for formulae C and D are as follows: the

15 subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11; the subscript 'h' is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8; the subscript 'i' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, and 8; the subscript 'j' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4;

20 and the subscript 'q' is an integer selected from the group consisting of 0, and 1. In one embodiment, X is F; Q is O; A is COOH and each R² is H.

Preparation of Linker Reagents and Support-Bound Activators

The polymer-supported perfluorosulfonyl fluoride linker of the present invention can be prepared as outlined in Figure 1.





Figure 1

Figure 1 illustrates the preparation of a linker **Figure 1, Structure 1**, **Structure 3** and a support-bound activator **Figure 1, Structure 4** from commercially available sulfonyl fluoride **Figure 1, Structure 1** (Aldrich Chemical Co., Milwaukee, Wisconsin, USA). Thus, treatment of iodide **Figure 1, Structure 1** with ethyl vinyl ether in the presence of sodium thiosulfate provides the aldehyde **Figure 1, Structure 2**. Oxidation of the aldehyde provides the carboxylic acid **Figure 1, Structure 3** which is suitable for attaching to a solid or semi-solid support via an amide- or ester-forming reaction with a suitable nucleophile on the support (e.g., an amino or hydroxy group). Alternatively, the carboxylic acid **Figure 1, Structure 3**, can be converted to its acid chloride using, for example, oxalyl chloride and then reacted with a suitable amine resin (e.g., Tentagel- NH_2 or Tentagel-NHEt). In still other embodiments, the aldehyde **Figure 1, Structure 2** can be attached to a resin via a reductive amination reaction of an amine resin, aldehyde **Figure 1, Structure 2** and a suitable reducing agent such as sodium borohydride.

Figure 2 illustrates the preparation of another linker reagent starting with 4-chloromethyl benzoyl chloride (Aldrich Chemical Co., Milwaukee, Wisconsin, USA).

Figure 2



5


Thus, treatment of 4-chloromethyl benzoyl chloride with t-butanol and triethylamine provides the ester **Figure 2, Structure 5** which can be converted to the thiol **Figure 2, Structure 6** upon treatment with thiourea, followed by hydrolysis with aqueous KOH. Oxidation of the thiol **Figure 2, Structure 6** with chlorine provides the sulfonyl chloride **Figure 2, Structure 7**. Conversion of the sulfonyl chloride to the sulfonate ester **Figure 2, Structure 8** can be accomplished with neopentyl alcohol in the presence of base (e.g., Et_3N). The sulfonate ester can then be activated by the stepwise exchange of the α hydrogen atoms with fluorine atoms ($t\text{BuLi}$ and NFSi) to give the ester **Figure 2, Structure 9**. Deprotection of the carboxylic ester with 20% TFA in methylene chloride provides the linker reagent **Figure 2, Structure 10**. The remaining steps in Figure 2 illustrate the attachment of the linker reagent to a support and the conversion to a sulfonic acid **Figure 2, Structure 12**.

Methods of Using the Support-Bound Activators

In yet another aspect, the present invention provides a method for covalently attaching a nucleophile to a compound having a hydroxy group or an enolizable ketone, the method comprising,

- 5 (a) contacting a compound having a hydroxy group or an enolizable ketone with a support-bound activator, wherein said contacting a compound having a hydroxy group or an enolizable ketone with a support bound activator forms an activated complex; and
- (b) contacting the activated complex with a reagent comprising a nucleophile under conditions sufficient to covalently attach the nucleophile to the compound.

10 In one embodiment, the support-bound activator has the formula:



wherein L is a linking group component; X is a member selected from the group consisting of F, and Cl; wherein the support-bound activator is covalently attached to a solid or semi-solid support.

- 15 Turning first to the solid or semi-solid support, the present invention is useful in a variety of solid-phase synthesis applications and, accordingly, a variety of supports find utility in this aspect of the invention. Typical solid supports include, but are not limited to, cross-linked divinylbenzene-styrene (polystyrene), controlled pore glass (CPG), polyacrylamides, poly(ethyleneglycol)monomethyl ether and poly(ethylene glycol) (PEG),
- 20 silica gel, cellulose, acrylic acid grafted polypropylene, and the like. Additionally, the solid support contains a reactive moiety suitable for attaching the linking group component. Suitably reactive moieties include, for example, a carboxylic acid, alcohol, amine, halomethyl and the like which is used to covalently attach the linking group component during construction of the present support-bound activators. Many of these
- 25 supports are available as functional polymers having reactive groups. Examples of such supports, include, by way of example, Acryloyl Wang resin, REM resin, Vinyl polystyrene, Vinylsulfonylmethyl polystyrene, (3-Formylindolyl)acetamidomethyl polystyrene, 2-(3,5-Dimethoxy-4-formylphenoxy)ethoxymethyl polystyrene, 2-(4-Formyl-3-methoxyphenoxy)ethyl polystyrene, 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin,
- 30 4-Benzyloxybenzaldehyde polystyrene, Aldehyde Wang resin, Formylpolystyrene, 1%

DVB, NovaSyn® TG acetal resin, Polystyrene-CHO, Carboxypolystyrene, NovaSyn® TG
 carboxy resin, Polystyrene-COOH, 4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-
 phenoxy resin, 4-Methylbenzhydramine resin HCl, 4-Methylbenzhydramine resin
 HCl, 9-Fmoc-amino-xanthen-3-yloxy, 9-Fmoc-amino-xanthen-3-yloxy, Amino-(4-
 5 methoxyphenyl)methyl polystyrene, Ethylamino-xanthen-3-yloxy-Merrifield resin,
 NovaSyn® TG Sieber resin, NovaSyn® TGR resin, Rink Amide AM resin, Rink Amide
 MBHA resin, Rink amide NovaGel™, Rink amide PEGA resin, Rink Amide resin, Sieber
 Amide resin, Sieber Ethylamide resin, Amino methyl resin, Amino PEGA resin,
 -Aminomethyl NovaGel™, -Aminomethylated polystyrene, N-Methylaminomethyl
 10 polystyrene, 4-Fmoc-hydrazinobenzoyl AM resin, 1H-Benzotriazole polystyrene,
 Benzotriazole-5-carbamidomethyl polystyrene, N-Fmoc- N-methoxy-β-alanine AM resin,
 Weinreb AM resin, 4-Sulfamylbenzoyl AM resin, (±)-1-(2,3-Isopropylidene)glycerol
 polystyrene, (±)-2,2-Dimethyldioxolan-4-methoxymethyl polystyrene, (±)-1-Glycerol
 polystyrene, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-
 15 Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-
 hydroxymethyl)-phenoxy resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid AM
 resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid BHA resin, 4-Hydroxymethyl-3-
 methoxyphenoxybutyric acid MBHA resin, 4-Hydroxymethylphenoxyacetyl NovaGel™,
 4-Hydroxymethylphenoxyacetyl PEGA resin, HMP resin, HMPA-NovaGel™, HMPB-
 20 AM resin, HMPB-BHA resin, HMPB-MBHA resin, Hydroxy-(2-chlorophenyl)methyl
 polystyrene, Hydroxymethylpolystyrene, NovaSyn® TG HMP resin, p-Benzyloxybenzyl
 Alcohol resin, Polystyrene-CH₂OH, Rink Acid resin, Trichloroacetimidate Wang resin,
 Wang resin, 4-Hydroxymethylbenzoic acid AM resin, 4-Hydroxymethylbenzoic acid
 NovaGel™, 4-Hydroxymethylbenzoic acid PEGA resin, 4-Hydroxyphenylsulfanylmethyl
 25 polystyrene, 9-(Hydroxymethyl)fluorene-4-carboxamidomethyl polystyrene, HESM
 polystyrene, HMBA-AM resin, HMBA-NovaGel™, HMBA-PEGA resin,
 Hydroxyethylsulfanylmethyl polystyrene, NovaSyn® TG HMBA resin, NovaSyn® TG
 hydroxy resin, Oxime resin, Aminoethyl photolinker resin, Hydroxyethyl photolinker
 resins, Hydroxymethyl photolinker resins, 3-[4-(Tritylmercapto)phenyl]propionyl AM
 30 resin, Mercaptomethyl polystyrene, NovaSyn® TG tritylthiol resin, Thiol 2-chlorotrityl
 resin, Thiol 4-methoxytrityl resin, (4-Bromophenyl)diisopropylsilyloxymethyl
 polystyrene, (4-Formylphenyl)diisopropylsilyloxymethyl polystyrene, (4-

Trityloxyphenyl)diisopropylsilyloxymethyl polystyrene. Solid supports also include TENTAGEL™, HYPOGEL™, JANDAJEL™, AND ARGOGEL™. Other solid supports include PEGylated polystyrene (polystyrene derivatized with polyethylene glycol), Tentagel-NH₂ resin, and derivatized Tentagel-NH₂ resin (e.g., by treatment with acetyl chloride followed by reduction with LiAlH₄ to provide Tentagel-NHEt resin). See also the Novabiochem Catalogue 2000 for additional resins and immobilized functional groups.



Such supports may take any size, shape or form, including particulate and non-particulate forms or shapes, spheres, disks, pellets, sheets, plugs, pins, crowns, lanterns, in beaded and non-beaded forms, resins, gels, microspheres, as well as amorphous forms and shapes. Embodiments of particulate supports, include beads, pellets, disks, amorphous particles, or other conventional forms. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-through devices. In a one embodiment, the diameter of the particulate support is 20-2000 micron, preferably 75-500 micron, more preferably 100-200 micron. As one skilled in the art will readily recognize, the scope of the present invention is not limited to the size, form, or shape of the solid or semi-solid support.

The linking group component can have a variety of structures. The linking group component is one which provides suitable spacing for the activator portion (-CF₂-SO₂-X) to interact freely with molecules or reactive components exposed to the activator portion. The linking group component is preferably 6-50 atoms long, more preferably 8-40 atoms long, even more preferably 8-30 atoms long, and yet more preferably 8-20 atoms long, thus providing sufficient exposure for the attached activator portion. Additionally, the linking group component, prior to attachment to the support, will have a attaching portion and a longer chain portion. The attaching portion is that part of the linking group component which can be directly attached to the solid support. This portion can be attached to the solid support via carbon-carbon bonds using, for example, supports having exposed (poly)trifluorochloroethylene moieties, or preferably, by siloxane bonds (using, for example, glass or silicon oxide as the solid support). Siloxane bonds the support are formed in one embodiment via reactions of attaching portions bearing trichlorosilyl or trialkoxysilyl groups. The attaching groups will also have a site for attachment of the longer chain portion. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl.

One skilled in the art will recognize that many additional methods of attaching linkers to solid and semi-solid supports exist. One method uses an amino resin to which an acid-bearing linker is attached via conventional techniques. Another method is the use of aryl ether linkages by coupling a phenol to a Merrifield (or equivalent) resin.

5 The longer chain portion can be any of a variety of molecules which are inert to the subsequent conditions used in the activator reactions described in further detail below. These longer chain portions can be ethylene glycol oligomers containing 2-14 monomer units, or more preferably 2-10 monomer units, and even more preferably 2-8 monomer units; in addition, the longer chain portions can be diamines, diacids, amino acids, peptides, or combinations thereof. In some embodiments, the longer chain portion also comprises an activator enhancing portion, i.e., a portion that increases the reactivity of the activator relative to an alkylene or ethylene glycol linking group. More particularly, an activator enhancing portion is one that provides additional electron withdrawing character to the activator portion (e.g., the $-\text{CF}_2-\text{SO}_2-\text{X}$ portion).

15 In one group of embodiments, X is a member selected from the group consisting of F and Cl. In another group of embodiments, L comprises an activator enhancing portion selected from the group consisting of:



20 wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, and NR^1 ; Z is a member selected from the group consisting of a chemical bond or CO; each R^2 is independently a member selected from the group consisting of hydrogen, halogen, (C_1-C_8) alkyl, (C_1-C_8) alkoxy, (C_2-C_8) heteroalkyl, (C_1-C_8) alkylthio, (C_1-C_8) alkylamino, $\text{di}(\text{C}_1-\text{C}_8)$ alkylamino, cyano, nitro and (C_1-C_8) alkylsulfonyl; the subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9,

and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).


In another embodiment, the support-bound activator is available in kit form for use in solid phase organic chemistry, as a reagent or catalyst in solution phase organic chemistry, as a scavenger resin in solution phase organic chemistry, as a silylating agent for use in analytical chemistry, and in particular, in chromatography, and as a reagent for the production of PET-ready molecules.

In one group of of embodiments, the support-bound activators of the present invention are:




15 wherein the symbol R^1 and the subscripts g, i, j, k and m all have the meanings provided above.

In another group of embodiments, the support-bound activators have a formula selected from:



and



20 In each of formulae A and B, X can be F and Cl; Q is O; Z is a chemical bond or C=O; Y is O-support or NR₁-support wherein R₁ is H, (C₁-C₈)alkyl or aryl and the support is a PEG-modified polystyrene or a Merrifield resin; and each R² is as defined more generally above. The subscripts in formulae A and B are as follows: 'g' is an integer
25 selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11; 'h' is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8; 'i' is an integer selected

from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, and 8; 'j' is an integer selected from the group consisting of 1, 2, 3, and 4; 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; 'm' is an integer selected from the group consisting of 2 and 3; 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4; and 'q' is an integer selected
5 from the group consisting of 1, and 2.

In one particular embodiment in formulae A and B, X is F; Q is O; Z is C=O, Y is NH-support wherein the support is a PEG-modified polystyrene; and each R² is H.

In one group of embodiments, the reagent is a member selected from the group consisting of an organostannane compound, an organozinc compound, an organoboron
10 compound, an organolithium compound, an organoaluminum compound, a Grignard reagent, an organosilicon compound, an organocopper compound, a thiol, a dialkylphosphite, an amine, a metal halide, and a halogen.

In one group of embodiments, the nucleophile is a member selected from the group consisting of an amine, a halogen anion, an aryl moiety, an alkyl moiety, a cyano, and a
15 hydride. In preferred embodiments, a transition metal catalyst is used in conjunction with the nucleophilic reagent.

In still other embodiments, the activated complex is treated under the appropriate conditions to afford a radioisotopically labeled compound without significant contamination from undesired compounds. This particular embodiment is expected to find
20 significant utility in the field of medicinal chemistry, in which the use of imaging agents remains an important technique to non-invasively evaluate human and animal conditions. One newly emerging field is that of positron emission tomography (PET). Briefly, this involves the use of short lived radioisotopes incorporated into known or potential drug substances and evaluation of their relative distribution throughout the body. One
25 significant drawback to existing methods is the preparation of the intended labeled compound and their use before the radioisotopes has decayed below that of useful intensity. One must prepare, purify, and use the intended labeled compound within a few hours in a PET study. Merely by way of example, U.S. Patent No. 6,307,372, entitled "Methods for high throughput chemical screening using magnetic resonance imaging" is
30 herein incorporated by reference in its entirety. Thus, there is a need in the art for a method for rapidly preparing and purifying PET-ready molecules. An approach that would

satisfy this need would be to provide support-bound targets that can be released from the support upon reaction with radioisotopically-labeled nucleophiles.

For example, treatment of a support-bound aryl perfluorosulfonate species with ^{11}C methyl lithium promotes a cleavage/derivatization cascade which results in the release of a ^{11}C methyl-substituted aryl species suitable for use in a PET experiment. Another example is incorporation of ^{18}F into aryl species to afford arylfluorides, commonly found in drug substances.

In one embodiment, the nucleophile is an ^{18}F anion. The labeled compounds resulting from the reaction of the ^{18}F anion with the activated complex can be used as PET-ready molecules for medical imaging purposes.

In still another embodiment, the nucleophile is a $^{11}\text{CH}_3$ anion. Examples of reagents for providing the $^{11}\text{CH}_3$ anion are ^{11}C methyl-lithium or ^{11}C methyl-cuprate. The labeled compounds resulting from the reaction of the $^{11}\text{CH}_3$ anion with the activated complex can be used as PET-ready molecules for medical imaging purposes.

In yet another embodiment, the present invention provides a method for transformation or substitution of a hydroxy group in a compound having a hydroxy group or an enolizable carbonyl group, the method comprising,

(a) contacting the compound having a hydroxy group or an enolizable carbonyl group with a support-bound activator having the formula:



wherein L is a linking group component; X is selected from F, and Cl; and the support bound activator is covalently attached to a solid or semi-solid support, under conditions sufficient to form an activated complex having the formula:



wherein -X is now -OQ and Q represents said compound; O is the oxygen atom vestige of the hydroxy group or enolizable ketone present in said compound; and L is the linking group component; and

(b) contacting the activated complex with a reagent comprising a nucleophile to transform or substitute said hydroxy group and form a new compound having the formula:



A variety of transformations can be accomplished using the methods herein. More particularly, the transformations include deoxygenations, cross-couplings (e.g., via organometallic reagents used in Suzuki, Stille, Heck, and organozinc coupling reactions), CO insertion, cyanide displacements, and numerous other reactions known to be suitable with vinyl or aryl triflates (see, Ritter, *Synthesis* 8:735-762 (1993)).

In one group of embodiments, the hydroxy group is present on an aromatic ring system such as a substituted or unsubstituted benzene or heteroaryl ring and the reagent is a hydride source which results in the reductive cleavage of the compound Q.

In one particular group of embodiments, the support-bound activator is a polymer-supported perfluorosulfonate linker as shown in Figure 3. Deoxygenation of a polymer-supported aryl perfluorosulfonate can be carried out by a palladium-mediated reduction. The polymer-bound aryl perfluorosulfonates can be efficiently cleaved with Et_3N/HCO_2H in the presence of a catalytic amount of $Pd(OAc)_2$ and 1,3-bis(diphenylphosphino)propane (dppp) in high yields to produce the reduced arenes under mild conditions. Table 1 provides examples of compounds (Q-OH) that can be reduced (Q-H) using the methods of the invention.

Figure 3









TABLE 1

	ArOH	ArH	Yield of ArH
14a			81
14b			76
14c			88
14d			90
14e			86
14f			85
14g			88
14h			75
14i			63
14j			

14k			86
14l			80

In another particular group of embodiments, the transformation is a cross-coupling reaction. In this group of embodiments, the activated complex is



5 wherein -X is now -OAr and Ar is an aryl or heteroaryl group.

In one group of embodiments, the reagent is an aryl boronic acid ($\text{Ar}'\text{B}(\text{OH})_2$) which provides a coupled compound having the formula $\text{Ar}-\text{Ar}'$. Figure 4 provides an illustration of this embodiment using a nonaflate linker of the present invention, and Table 2 provides an illustration of the compounds and products provided. For simplicity, the linker is not shown in Table 2.

Figure 4


























TABLE 2

Resin	Aryl boronic acid	Product (Ar'-Ar)
		 16a
		 16b
		 16c
		 16d
		 16e
		 16f
		 16g



In still other embodiments, the activated complex is subjected to additional synthetic transformations prior to a cross-coupling (e.g., Suzuki, Stille, Heck, Sonogashira, Buchwald) which releases the product from the resin. Figure 5 illustrates certain reactions that can be applied to the activated complex, while Table 3 illustrates various starting materials and products that can be obtained using the linking group/activators described herein.




























TABLE 3


R^1NH_2	R^2COCl	Boronic Acid	Product
			 20a
			 20b
			 20c
			 20d
			 20e
			 20f


20g			
20h			
20i			

The perfluorosulfonyl fluoride linker of this invention was used to a multi-step synthesis of the therapeutic agent, meclizine as outlined in Figure 6.

5


Figure 6





5 *Strongly Acidic Support*

In another aspect, the present invention provides a strongly acidic support comprising a solid or semi-solid support; and at least one support-bound strongly acid group having the formula:



- 10 wherein L is a linking group component; and X is OH; and wherein the support-bound strongly acid group is covalently attached to the solid or semi-solid support.

Turning first to the solid or semi-solid support, the present invention is useful in a variety of solid-phase synthesis applications and, accordingly, a variety of supports find utility in this aspect of the invention. Typical solid supports include, but are not limited to,

15 cross-linked divinylbenzene-styrene (polystyrene), controlled pore glass (CPG), polyacrylamides, poly(ethyleneglycol)monomethyl ether and poly(ethylene glycol) (PEG), silica gel, cellulose, acrylic acid grafted polypropylene, and the like. Additionally, the solid support contains a reactive moiety suitable for attaching the linking group component. Suitably reactive moieties include, for example, a carboxylic acid, alcohol,

20 amine, halomethyl and the like which is used to covalently attach the linking group component during construction of the present support-bound activators. Many of these supports are available as functional polymers having reactive groups. Examples of such

supports, include, by way of example, Acryloyl Wang resin, REM resin, Vinyl
 polystyrene, Vinylsulfonylmethyl polystyrene, (3-Formylindolyl)acetamidomethyl
 polystyrene, 2-(3,5-Dimethoxy-4-formylphenoxy)ethoxymethyl polystyrene, 2-(4-Formyl-
 3-methoxyphenoxy)ethyl polystyrene, 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin,
 5 4-Benzyloxybenzaldehyde polystyrene, Aldehyde Wang resin, Formylpolystyrene, 1%
 DVB, NovaSyn® TG acetal resin, Polystyrene-CHO, Carboxypolystyrene, NovaSyn® TG
 carboxy resin, Polystyrene-COOH, 4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-
 phenoxy resin, 4-Methylbenzhydramine resin HCl, 4-Methylbenzhydramine resin
 HCl, 9-Fmoc-amino-xanthen-3-yloxy, 9-Fmoc-amino-xanthen-3-yloxy, Amino-(4-
 10 methoxyphenyl)methyl polystyrene, Ethylamino-xanthen-3-yloxy-Merrifield resin,
 NovaSyn® TG Sieber resin, NovaSyn® TGR resin, Rink Amide AM resin, Rink Amide
 MBHA resin, Rink amide NovaGel™, Rink amide PEGA resin, Rink Amide resin, Sieber
 Amide resin, Sieber Ethylamide resin, Amino methyl resin, Amino PEGA resin,
 Aminomethyl NovaGel™, Aminomethylated polystyrene, N-Methylaminomethyl
 15 polystyrene, 4-Fmoc-hydrazinobenzoyl AM resin, 1H-Benzotriazole polystyrene,
 Benzotriazole-5-carbamidomethyl polystyrene, N-Fmoc- N-methoxy-β-alanine AM resin,
 Weinreb AM resin, 4-Sulfamylbenzoyl AM resin, (±)-1-(2,3-Isopropylidene)glycerol
 polystyrene, (±)-2,2-Dimethyldioxolan-4-methoxymethyl polystyrene, (±)-1-Glycerol
 polystyrene, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-
 20 Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-
 hydroxymethyl)-phenoxy resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid AM
 resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid BHA resin, 4-Hydroxymethyl-3-
 methoxyphenoxybutyric acid MBHA resin, 4-Hydroxymethylphenoxyacetyl NovaGel™,
 4-Hydroxymethylphenoxyacetyl PEGA resin, HMP resin, HMPA-NovaGel™, HMPB-
 25 AM resin, HMPB-BHA resin, HMPB-MBHA resin, Hydroxy-(2-chlorophenyl)methyl
 polystyrene, Hydroxymethylpolystyrene, NovaSyn® TG HMP resin, p-Benzyloxybenzyl
 Alcohol resin, Polystyrene-CH₂OH, Rink Acid resin, Trichloroacetimidate Wang resin,
 Wang resin, 4-Hydroxymethylbenzoic acid AM resin, 4-Hydroxymethylbenzoic acid
 NovaGel™, 4-Hydroxymethylbenzoic acid PEGA resin, 4-Hydroxyphenylsulfanylmethyl
 30 polystyrene, 9-(Hydroxymethyl)fluorene-4-carboxamidomethyl polystyrene, HESM
 polystyrene, HMBA-AM resin, HMBA-NovaGel™, HMBA-PEGA resin,
 Hydroxyethylsulfanylmethyl polystyrene, NovaSyn® TG HMBA resin, NovaSyn® TG

hydroxy resin, Oxime resin, Aminoethyl photolinker resin, Hydroxyethyl photolinker resins, Hydroxymethyl photolinker resins, 3-[4-(Tritylmercapto)phenyl]propionyl AM resin, Mercaptomethyl polystyrene, NovaSyn® TG tritylthiol resin, Thiol 2-chlorotrityl resin, Thiol 4-methoxytrityl resin, (4-Bromophenyl)diisopropylsilyloxymethyl polystyrene, (4-Formylphenyl)diisopropylsilyloxymethyl polystyrene, (4-Trityloxyphenyl)diisopropylsilyloxymethyl polystyrene. Solid supports also include TENTAGEL™, HYPOGEL™, JANDAJEL™, AND ARGOGEL™. Other solid supports include PEGylated polystyrene (polystyrene derivatized with polyethylene glycol), Tentagel-NH₂ resin, and derivatized Tentagel-NH₂ resin (e.g., by treatment with acetyl chloride followed by reduction with LiAlH₄ to provide Tentagel-NHEt resin). See also the Novabiochem Catalogue 2000 for additional resins and immobilized functional groups.

Such supports may take any size, shape or form, including particulate and non-particulate forms or shapes, spheres, disks, pellets, sheets, plugs, pins, crowns, lanterns, in beaded and non-beaded forms, resins, gels, microspheres, as well as amorphous forms and shapes. Embodiments of particulate supports, include beads, pellets, disks, amorphous particles, or other conventional forms. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-through devices. In a one embodiment, the diameter of the particulate support is 20-2000 micron, preferably 75-500 micron, more preferably 100-200 micron. As one skilled in the art will readily recognize, the scope of the present invention is not limited to the size, form, or shape of the solid or semi-solid support.


The linking group component can have a variety of structures. The linking group component is one which provides suitable spacing for the activator portion (-CF₂-SO₂-OH) to interact freely with molecules or reactive components exposed to the activator portion. The linking group component is preferably 6-50 atoms long, more preferably 8-40 atoms long, even more preferably 8-30 atoms long, and yet more preferably 8-20 atoms long, thus providing sufficient exposure for the attached activator portion. Additionally, the linking group component, prior to attachment to the support, will have a attaching portion and a longer chain portion. The attaching portion is that part of the linking group component which can be directly attached to the solid support. This portion can be attached to the solid support via carbon-carbon bonds using, for example, supports having exposed (poly)trifluorochloroethylene moieties, or preferably, by siloxane bonds (using,

for example, glass or silicon oxide as the solid support). Siloxane bonds the support are formed in one embodiment via reactions of attaching portions bearing trichlorosilyl or trialkoxysilyl groups. The attaching groups will also have a site for attachment of the longer chain portion. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl.

One skilled in the art will recognize that many additional methods of attaching linkers to solid or semi-solid supports exist. One method uses an amino resin to which an acid-bearing linker is attached via conventional techniques. Another method is the use of aryl ether linkages by coupling a phenol to a Merrifield (or equivalent) resin.

10 The longer chain portion can be any of a variety of molecules which are inert to the subsequent conditions used in the activator reactions described in further detail below. These longer chain portions can be ethylene glycol oligomers containing 2-14 monomer units, or more preferably 2-10 monomer units, and even more preferably 2-8 monomer units; in addition, the longer chain portions can be diamines, diacids, amino acids, 15 peptides, or combinations thereof. In some embodiments, the longer chain portion also comprises an activator enhancing portion, i.e., a portion that increases the reactivity of the activator relative to an alkylene or ethylene glycol linking group. More particularly, an activator enhancing portion is one that provides additional electron withdrawing character to the activator portion (e.g., the $-\text{CF}_2-\text{SO}_2-\text{OH}$ portion).


20 In another group of embodiments, L comprises an activator enhancing portion
selected from the group consisting of:




wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, and NR¹; Z is a member selected from the group consisting of a chemical bond or CO; each R² is independently a member selected from the group consisting of hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-C₈)alkylthio, (C₁-

C_8)alkylamino, di(C_1 - C_8)alkylamino, cyano, nitro and (C_1 - C_8)alkylsulfonyl; the subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).

In another group of embodiments, the support-bound activators have a formula selected from:



and



In each of formulae A and B, X is OH; Q is O; Z is a chemical bond or C=O; Y is O-support or NR_1 -support wherein R_1 is H, (C_1 - C_8)alkyl or aryl and the support is a PEG-modified polystyrene or a Merrifield resin; and each R^2 is as defined more generally above. The subscripts in formulae A and B are as follows: 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11; 'h' is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8; 'i' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, and 8; 'j' is an integer selected from the group consisting of 1, 2, 3, and 4; 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; 'm' is an integer selected from the group consisting of 2 and 3; 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4; and 'q' is an integer selected from the group consisting of 1, and 2.

In one particular embodiment in formulae A and B, X is OH; Q is O; Z is C=O, Y is NH-support wherein the support is a PEG-modified polystyrene; and each R² is H.

In still other embodiments, each solid or semi-solid support comprises a plurality of support-bound strongly acid groups, with a density of at least 1 nmol support-bound strongly acid groups per gram of solid or semi-solid support, more preferably, at least 1 μmol support-bound strongly acid groups per gram of solid or semi-solid support, and even more preferably, at least 1 mmol support-bound strongly acid groups per gram of solid or semi-solid support.

In another embodiment, the strongly acidic support is available in kit form for use as a reagent or catalyst in solution phase organic chemistry, or as a scavenger resin in solution phase organic chemistry.

The preparation of the resin-bound perfluorosulfonic acid **Figure 7, Structure 25** is outlined in Figure 7, and the synthesis of the resin-bound perfluorosulfonic acid **Figure 2, Structure 12** is outlined Figure 2.




Figure 7

The highly acidic supports of this invention provide a solid phase reagent that is chemically analogous to perfluorosulfonic acids; furthermore, in one embodiment of the present invention, the vast majority of surface-bound perfluorosulfonic acid sites are readily available for reaction, thus allowing the use of this embodiment both catalytically and stoichiometrically.

In addition to being used as highly acidic solid phase catalysts and reagents, support-bound acidic groups can be used as scavenger resins. Such a use has been described in the generation of diverse collections of compounds. Merely by way of

example, "Polymer Supported Reagents Handbook" (NovaBiochem, 2001), is incorporated herein by reference. In general, resins capable of capturing excess reagents, products, or unwanted byproducts from reaction mixtures have found widespread use in organic chemistry laboratories, and in particular, are considered to be highly important in the pharmaceutical arena. These resins can be employed in both covalent and ionic fashions.

For example, an amine can react with an acid chloride to afford an amide as the desired product. The most common method of obtaining the product in a high yield is to use one of the two reaction partners in excess and to then separate the product from excess reagent. When using an excess of the acid chloride, the reaction mixture can be treated with a primary or secondary amine-containing resin after the reaction to *covalently* capture the excess acid chloride as a polymer bound amide. Simple filtration of the resin from the reaction mixture would afford the desired product amide essentially free of excess acid chloride. Alternatively, when using an excess of the amine component, the reaction mixture can be treated with an acid-containing resin (i.e., a strong cation exchange resin) to *ionically* capture the excess amine as its ammonium salt form on the resin. Simple filtration of the resin from the reaction mixture would afford the desired amide essentially free of the excess amine.

To date there have been no scavenger resins described containing acidic groups stronger than phenylsulfonic acids (e.g., DOWEX™ (Dow Chemical Company, Midland, Michigan) and related resins). These resins are inherently limited in their ability to capture weakly basic compounds by virtue of their modest acidity. There exists the need to develop resins of much higher acidity (i.e. highly acidic supports) which are swellable in a variety of solvents, capable of achieving high loading (i.e. loading of > 0.2 mmol/g, preferably > 0.5 mmol/g, more preferably > 0.8 mmol/g), and stable to mechanical forces typical during washing steps.

Another application of scavenger supports has been termed "capture and release," a technique in which targets are first captured (i.e., attached to a support) from a chemical reaction solution, then, the unwanted compounds removed by filtration or other similar methods, and finally, the desired compound is released from the support. See Bhat, *J. Comb. Chem.* 2000, 2, 597. This final release step can be performed by a multitude of possible methods depending on the nature of the attachment of the targets to the support.

When using resins to "capture" molecules to the support as salts, the displacement step may be conveniently performed by treating the support with a solution containing a volatile amine such as ammonia. Molecules covalently attached to the support can be cleaved under a variety of conditions.

5 Thus, another embodiment of the polymer-supported compositions described in this is the use of the polymer-supported compositions as either covalent or ionic scavenger resins. An embodiment of this invention is to capture amines, thiols, alcohols, phenols, ketones and other species capable of forming stable triflate products as their respective polymer-supported triflates, and then to filter them away from reaction mixtures. A
10 preferred embodiment of this invention is to expose a reaction mixture containing amine and other basic compounds capable of forming ionic triflate salts with the novel compositions disclosed herein and to then filter away said basic species. See Example 8 below for an illustrative use of the highly acidic supports of the present invention as scavenger resins.

15 Those skilled in the art will recognize that the polymer-supported perfluorsulfonic acids described herein are dramatically more acidic than the known cation exchange resins, and as such, will be capable of effectively scavenging a broader range of compounds.


20 *Silylating Support*

 In the field of analytical chemistry, it is common to chemically derivatize molecules to make them more suitable for various chromatographic techniques. The general principles of compound derivatization for chromatography are described in
25 "Handbook of Derivatives for Chromatography: by K. Blau and J. Halket (Wiley, England, 1993), which is incorporated herein by reference. For example, treatment of molecules containing hydroxyl or acid groups with any one of a number of silylating agents renders silyl ethers or esters that are more volatile than the parent compound, and thus more amenable to analysis via gas chromatography. It is also common to derivatize
30 compounds to alter their properties in HPLC applications. One drawback in current methods for compound derivatization is the necessity to separate excess derivatization reagents from the desired, derivatized product. In addition, highly reactive derivatization

regents, such as chlorotrimethylsilane and other silylating reagents, have been observed to both deactivate chromatography columns and contaminate flame ionization detectors in gas chromatographs with SiO₂ rendering them ineffective. A technique of employing highly reactive, polymer-supported derivatization reagents would be very important and find widespread use through the industry.

One aspect of the present invention provides for a silylating support, and in particular, silylating polymer-supported reagents, that afford the desired derivatized products free of contaminating reagents. Whereas silyltriflates are widely known as extremely reactive silylating reagents in solution, the corresponding polymer-supported silyltriflates have not been widely employed. NAFION™-TMS has been made by Noyori, *Tetrahedron Lett.* **1980**, *21*, 767 by heating the acid form of the polymer with chlorotrimethylsilane and sulfuric acid. Furthermore, the polymer-supported silyl triflates that have been described in the literature (e.g., Smith, *Tetrahedron Lett.* **1999**, *40*, 3285 and Porco, *Tetrahedron Lett.* **1999**, *40*, 3289) are trifylating resins and not silylating resins. Thus, there is a need in the art for highly-reactive, silylating resins that are swellable and wettable by organic solvents, and which can thus readily act as stoichiometric silylating reagents. The silylating support of the present invention fulfills this need.

In another aspect, the present invention provides a silylating support comprising a solid or semi-solid support; and at least one support-bound silylating group having the formula:



wherein L is a linking group component; X is a trisubstituted silyloxy; wherein the support-bound silylating group is covalently attached to the solid or semi-solid support.

Turning first to the solid or semi-solid support, the present invention is useful in a variety of solid-phase synthesis applications and, accordingly, a variety of supports find utility in this aspect of the invention. Typical solid supports include, but are not limited to, cross-linked divinylbenzene-styrene (polystyrene), controlled pore glass (CPG), polyacrylamides, poly(ethyleneglycol)monomethyl ether and poly(ethylene glycol) (PEG), silica gel, cellulose, acrylic acid grafted polypropylene, and the like. Additionally, the

solid support contains a reactive moiety suitable for attaching the linking group component. Suitably reactive moieties include, for example, a carboxylic acid, alcohol, amine, halomethyl and the like which is used to covalently attach the linking group component during construction of the present support-bound activators. Many of these

5 supports are available as functional polymers having reactive groups. Examples of such supports, include, by way of example, Acryloyl Wang resin, REM resin, Vinyl polystyrene, Vinylsulfonylmethyl polystyrene, (3-Formylindolyl)acetamidomethyl polystyrene, 2-(3,5-Dimethoxy-4-formylphenoxy)ethoxymethyl polystyrene, 2-(4-Formyl-3-methoxyphenoxy)ethyl polystyrene, 4-(4-Formyl-3-methoxyphenoxy)butyryl AM resin,

10 4-Benzyloxybenzaldehyde polystyrene, Aldehyde Wang resin, Formylpolystyrene, 1% DVB, NovaSyn® TG acetal resin, Polystyrene-CHO, Carboxypolystyrene, NovaSyn® TG carboxy resin, Polystyrene-COOH, 4-(2',4'-Dimethoxyphenyl-Fmoc-aminomethyl)-phenoxy resin, 4-Methylbenzhydramine resin HCl, 4-Methylbenzhydramine resin HCl, 9-Fmoc-amino-xanthen-3-yloxy, 9-Fmoc-amino-xanthen-3-yloxy, Amino-(4-

15 methoxyphenyl)methyl polystyrene, Ethylamino-xanthen-3-yloxy-Merrifield resin, NovaSyn® TG Sieber resin, NovaSyn® TGR resin, Rink Amide AM resin, Rink Amide MBHA resin, Rink amide NovaGel™, Rink amide PEGA resin, Rink Amide resin, Sieber Amide resin, Sieber Ethylamide resin, Amino methyl resin, Amino PEGA resin, Aminomethyl NovaGel™, Aminomethylated polystyrene, N-Methylaminomethyl

20 polystyrene, 4-Fmoc-hydrazinobenzoyl AM resin, 1H-Benzotriazole polystyrene, Benzotriazole-5-carbamidomethyl polystyrene, N-Fmoc- N-methoxy-β-alanine AM resin, Weinreb AM resin, 4-Sulfamylbenzoyl AM resin, (±)-1-(2,3-Isopropylidene)glycerol polystyrene, (±)-2,2-Dimethyldioxolan-4-methoxymethyl polystyrene, (±)-1-Glycerol polystyrene, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-

25 Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-(2',4'-Dimethoxyphenyl-hydroxymethyl)-phenoxy resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid AM resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid BHA resin, 4-Hydroxymethyl-3-methoxyphenoxybutyric acid MBHA resin, 4-Hydroxymethylphenoxyacetyl NovaGel™, 4-Hydroxymethylphenoxyacetyl PEGA resin, HMP resin, HMPA-NovaGel™, HMPB-

30 AM resin, HMPB-BHA resin, HMPB-MBHA resin, Hydroxy-(2-chlorophenyl)methyl polystyrene, Hydroxymethylpolystyrene, NovaSyn® TG HMP resin, p-Benzyloxybenzyl Alcohol resin, Polystyrene-CH₂OH, Rink Acid resin, Trichloroacetimidate Wang resin,

Wang resin, 4-Hydroxymethylbenzoic acid AM resin, 4-Hydroxymethylbenzoic acid NovaGel™, 4-Hydroxymethylbenzoic acid PEGA resin, 4-Hydroxyphenylsulfanylmethyl polystyrene, 9-(Hydroxymethyl)fluorene-4-carboxamidomethyl polystyrene, HESM polystyrene, HMBA-AM resin, HMBA-NovaGel™, HMBA-PEGA resin, 5 Hydroxyethylsulfanylmethyl polystyrene, NovaSyn® TG HMBA resin, NovaSyn® TG hydroxy resin, Oxime resin, Aminoethyl photolinker resin, Hydroxyethyl photolinker resins, Hydroxymethyl photolinker resins, 3-[4-(Tritylmercapto)phenyl]propionyl AM resin, Mercaptomethyl polystyrene, NovaSyn® TG tritylthiol resin, Thiol 2-chlorotrityl resin, Thiol 4-methoxytrityl resin, (4-Bromophenyl)diisopropylsilyloxymethyl 10 polystyrene, (4-Formylphenyl)diisopropylsilyloxymethyl polystyrene, (4-Trityloxyphenyl)diisopropylsilyloxymethyl polystyrene. Solid supports also include TENTAGEL™, HYPOGEL™, JANDAJEL™, AND ARGOGEL™. Other solid supports include PEGylated polystyrene (polystyrene derivatized with polyethylene glycol), Tentagel-NH₂ resin, and derivatized Tentagel-NH₂ resin (e.g., by treatment with acetyl 15 chloride followed by reduction with LiAlH₄ to provide Tentagel-NHEt resin). See also the Novabiochem Catalogue 2000 for additional resins and immobilized functional groups.

Such supports may take any size, shape or form, including particulate and non-particulate forms or shapes, spheres, disks, pellets, sheets, plugs, pins, crowns, lanterns, in beaded and non-beaded forms, resins, gels, microspheres, as well as amorphous forms and 20 shapes. Embodiments of particulate supports, include beads, pellets, disks, amorphous particles, or other conventional forms. The solid or semi-solid supports may be used as single particle, as groups of particles, as free flowing particles, and may be packed into columns, tubes or other flow-through devices. In a one embodiment, the diameter of the particulate support is 20-2000 micron, preferably 75-500 micron, more preferably 100-200 25 micron. As one skilled in the art will readily recognize, the scope of the present invention is not limited to the size, form, or shape of the solid or semi-solid support.



The linking group component can have a variety of structures. The linking group component is one which provides suitable spacing for the activator portion (-CF₂-SO₂-OX; wherein X is a trisubstituted silyloxy) to interact freely with molecules or reactive 30 components exposed to the activator portion. The linking group component is preferably 6-50 atoms long, more preferably 8-40 atoms long, even more preferably 8-30 atoms long, and yet more preferably 8-20 atoms long, thus providing sufficient exposure for the

attached activator portion. Additionally, the linking group component, prior to attachment to the support, will have a attaching portion and a longer chain portion. The attaching portion is that part of the linking group component which can be directly attached to the solid support. This portion can be attached to the solid support via carbon-carbon bonds
5 using, for example, supports having exposed (poly)trifluorochloroethylene moieties, or preferably, by siloxane bonds (using, for example, glass or silicon oxide as the solid support). Siloxane bonds the support are formed in one embodiment via reactions of attaching portions bearing trichlorosilyl or trialkoxysilyl groups. The attaching groups will also have a site for attachment of the longer chain portion. For example, groups
10 which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl.

One skilled in the art will recognize that many additional methods of attaching linkers to solid or semi-solid supports exist. One method uses an amino resin to which an acid-bearing linker is attached via conventional techniques. Another method is the use of
15 aryl ether linkages by coupling a phenol to a Merrifield (or equivalent) resin.


The longer chain portion can be any of a variety of molecules which are inert to the subsequent conditions used in the activator reactions described in further detail below. These longer chain portions can be ethylene glycol oligomers containing 2-14 monomer units, or more preferably 2-10 monomer units, and even more preferably 2-8 monomer
20 units; in addition, the longer chain portions can be diamines, diacids, amino acids, peptides, or combinations thereof. In some embodiments, the longer chain portion also comprises an activator enhancing portion, i.e., a portion that increases the reactivity of the activator relative to an alkylene or ethylene glycol linking group. More particularly, an activator enhancing portion is one that provides additional electron withdrawing character
25 to the activator portion (e.g., the $-\text{CF}_2-\text{SO}_2-\text{OX}$ portion; wherein X is a trisubstituted silyloxy).

In another group of embodiments, L comprises an activator enhancing portion selected from the group consisting of:




- 5 wherein Y is a member selected from the group consisting of a chemical bond, O, CO, S, and NR^1 ; Z is a member selected from the group consisting of a chemical bond or CO; each R^2 is independently a member selected from the group consisting of hydrogen, halogen, $(\text{C}_1\text{-C}_8)\text{alkyl}$, $(\text{C}_1\text{-C}_8)\text{alkoxy}$, $(\text{C}_2\text{-C}_8)\text{heteroalkyl}$, $(\text{C}_1\text{-C}_8)\text{alkylthio}$, $(\text{C}_1\text{-C}_8)\text{alkylamino}$, $\text{di}(\text{C}_1\text{-C}_8)\text{alkylamino}$, cyano, nitro and $(\text{C}_1\text{-C}_8)\text{alkylsulfonyl}$; the subscript 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'i' is an integer selected from the group consisting of 1, 2, 3, 4, 5, and 6; the subscript 'j' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10; the subscript 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; the subscript 'm' is an integer selected from the group consisting of 2 and 3; and the subscript 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4. In the groups described herein as dialkylamino, the alkyl groups can be the same or different, or can optionally be combined to form a ring having additional heteroatoms (e.g., pyrrolidino, morpholino, piperazino).

- 20 In another group of embodiments, the support-bound activators have a formula selected from:



and



In each of formulae A and B, X is trisubstituted silyloxy; Q is O; Z is a chemical bond or C=O; Y is O-support or NR₁-support wherein R₁ is H, (C₁-C₈)alkyl or aryl and the support is a PEG-modified polystyrene or a Merrifield resin; and each R² is as defined more generally above. The subscripts in formulae A and B are as follows: 'g' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11; 'h' is an integer selected from the group consisting of 1, 2, 3, 4, 5, 6, 7, and 8; 'i' is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6, 7, and 8; 'j' is an integer selected from the group consisting of 1, 2, 3, and 4; 'k' is an integer selected from the group consisting of 1, 2, 3, and 4; 'm' is an integer selected from the group consisting of 2 and 3; 'n' is an integer selected from the group consisting of 0, 1, 2, 3, and 4; and 'q' is an integer selected from the group consisting of 1, and 2.

In one particular embodiment in formulae A and B, X is trisubstituted silyloxy; Q is O; Z is C=O, Y is NH-support wherein the support is a PEG-modified polystyrene; and each R² is H.

In one embodiment, the trisubstituted silyloxy has the formula OSiR³R⁴R⁵, wherein, each of R³, R⁴, and R⁵ is independently a member selected from the group consisting of substituted (C₁-C₈)alkyl, unsubstituted (C₁-C₈)alkyl, substituted (C₁-C₈)alkenyl, unsubstituted (C₁-C₈)alkenyl, substituted aryl, and unsubstituted aryl.

In another embodiment, each of R³, R⁴, R⁵ is a methyl group; in another embodiment, each of R³, R⁴ is a methyl group and R⁵ is a t-butyl group.

In still other embodiments, each solid or semi-solid support comprises a plurality of support-bound silylating groups, with a concentration of at least 1nmol support-bound silylating groups per gram of solid or semi-solid support, more preferably, at least 1μmol support-bound silylating groups per gram of solid or semi-solid support, and even more preferably, at least 1 mmol support-bound silylating groups per gram of solid or semi-solid support.

In another embodiment, the silylating support of the present invention can be provided in kit form for pre-treatment of test compounds prior to analysis by a wide range of analytical chemistry techniques, including gas chromatography and high-performance liquid chromatography. Such a kit could provide the silylating support in the form of a resin or particles packed into a column. The compound to be silylated could then be passed over or through the silylating support prior to analysis by chromatography.

See Example 9 below for one illustrative use of the silylating supports of the present invention.

The following examples provide more detailed descriptions of synthetic methods used to prepare traceless linkers and activators of the present invention. One of skill in the art will appreciate that many of the methods provided below can be applied to the preparation of other linkers and activators. Accordingly, the examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1

This example illustrates the preparation of a polymer-supported perfluorosulfonyl fluoride linker (Figure 1).

At 0 °C, to a stirring solution containing ethyl vinyl ether (600 mg, 8.3 mmol), NaHCO₃ (680 mg, 8.0 mmol), and commercial tetrafluoro-2-(tetrafluoro-2-iodoethoxy)ethanesulfonyl fluoride **Figure 1, Structure 1** (3.5 g, 8.0 mmol) in CH₃CN (8 mL) and H₂O (7 mL) was slowly added Na₂S₂O₄ (1.4 g, 8.0 mmol). The reaction mixture was stirred at 5 °C for 50 min. The pH of the reaction mixture was adjusted to 6.2~7.0 by adding 3.0 N aqueous HCl and the mixture was stirred at 25 °C for another 20 min. The reaction mixture was extracted with CH₂Cl₂, washed with water and concentrated under reduced pressure. The oily residue was dissolved in acetone (38 mL) and the solution was added to a stirring mixture of 2-methyl-butene-2 (36 mL), NaH₂PO₄ (4.0 g, mmol), NaClO₂ (5.0 g, mmol) and water (40 mL) at 0-5 °C. The reaction mixture was stirred at 5-15 °C for 2 h. The reaction mixture was concentrated under reduced pressure, extracted with ether, washed with brine, dried (MgSO₄) and concentrated. The oily crude product was purified on a SiO₂ column (MeOH/CHCl₃, 2:98) to give 1.8 g (62%) of the linker **Figure 1, Structure 3** as thick oil. ¹H NMR (CDCl₃, δ): 3.17 (t, J=16.8 Hz, CF₂CH₂CO₂H); ¹³C NMR (CDCl₃, δ): 169.54, 36.69, 36.39, 36.09; ¹⁹F NMR (CDCl₃, δ): 121.61 (SO₂F), 6.08, 11.51, -36.14, -39.76; MS (ESI) calcd for C₆H₃F₉O₅S 357.96, found: 357.0 (M-H)⁻.

To a stirring solution of the above acid linker **Figure 1, Structure 3** (800 mg, 2.24 mmol) and oxalyl chloride (430 μ L, 4.8 mmol) in CH_2Cl_2 (1.6 mL) was slowly added DMF (18 μ L). After the evolution of gas bubbles, the reaction mixture was stirred for another 1 h and then concentrated under reduced pressure. The oily product was dissolved in CH_2Cl_2 (5 mL), the solution was added to a commercial TENTAGEL™ NH_2 resin (1.08 g, 0.46 mmol) and the resin was cooled in a dry-ice container for 10 min. To the resin was slowly added diisopropylethylamine (1.2 mL, 7.0 mmol), and the resin was shaken at room temperature overnight. The beads were washed with CH_2Cl_2 and dried under vacuum overnight to give resin-bound linker **Figure 1, Structure 4a**.

10

Example 2

This example illustrates the preparation of difluorosulfonic acid linker and the resin-bound difluorosulfonic acid (Figure 2).

A solution of chloride **Figure 2, Structure 5** (3.8 g, 16.8 mmol) and thiourea (1.3 g, 17 mmol) in EtOH (10 mL) was stirred at 70 °C for 4 h and then cooled to room temperature. A solution of NaOH (1.2 g, 30 mmol) in water (10 mL) was added, and the reaction mixture was stirred at room temperature overnight, concentrated under reduced pressure, acidified to pH 6, extracted with CH_2Cl_2 , washed with water. The crude thiol **Figure 2, Structure 6** was dissolved in CH_2Cl_2 (30 mL), water (30 mL) and acetic acid (2 mL). The solution was cooled to 0-5 °C and Cl_2 gas was bubbled onto the solution for 1 h. The reaction mixture was concentrated, extracted with CH_2Cl_2 , washed with cold water, dried (MgSO_4) and concentrated to give sulfonyl chloride **Figure 2, 7** (4.8 g, 97%). ^1H NMR (CDCl_3 , δ): 1.60 (s, 9 H), 4.92 (s, 2 H), 7.68 (d, 2 H), 8.38 (d, 2 H); ^{13}C NMR (CDCl_3 , δ): 28.34, 70.52, 81.92, 130.37, 131.44, 134.02, 164.98.

Et_3N (2.0 g, 19.5 mmol) was added to a solution of **Figure 2, 7** (4.2 g, 14 mmol), neopentyl alcohol (2.6 g, 30 mmol) in CH_2Cl_2 (20 mL) at -78 °C. The reaction mixture was stirred at room temperature overnight, diluted with cold water, extracted with CH_2Cl_2 , washed with water, dried (MgSO_4) and concentrated under reduced pressure to give crude product **Figure 2, 8** as light yellow solid. The crude product was washed with a mixture of hexane and ether (4:1) to yield pure **Figure 2, 8** as off-white powder (4.3 g, 90%). ^1H NMR (CDCl_3 , δ): 0.96 (s, 9 H), 1.60 (s, 9 H), 3.78 (s, 2 H), 4.40 (s, 2 H), 7.44 (d, 2 H),

8.02 (d, 2 H); ^{13}C NMR (CDCl_3 , δ): 26.18, 28.36, 32.03, 56.50, 79.83, 81.64, 130.07, 130.74, 132.48, 132.87, 165.29.

To a solution of the benzylic sulfonate ester **Figure 2, 8** (342 mg, 1.0 mmol) in anhydrous THF at -78°C was added *t*-BuLi (620 μL , 1.05 mmol) over a period of 2 min, and the mixture was stirred for 30 min at -78°C . A solution of NFSi (341 mg, 1.05 mmol) in THF (0.5 mL) was added dropwise at -78°C . The mixture was stirred at -78°C for 1 h. The procedure was repeated with *t*-BuLi (650 μL , 1.10 mmol) and NFSi (357 mg, 1.10 mmol). The reaction was quenched with water, extracted with CHCl_3 , washed with water, dried (MgSO_4) and concentrated. The crude product was purified on a SiO_2 column (MeOH/ CHCl_3 , 2:98) to give pure **Figure 2, 9**. A solution of trifluoroacetic acid (1.8 mL) in CH_2Cl_2 (4 mL) was added to **Figure 2, 9** and the mixture was stirred for 40 min, concentrated to give **Figure 2, 10** as a white powder (252 mg, 78% from 9). ^1H NMR (CD_3COCD_3 , δ): 1.02 (s, 9 H), 4.22 (s, 2 H), 7.91 (d, 2 H), 8.26 (d, 2 H).

Et_3N (30 mg, 0.30 mmol) was added to a solution of **Figure 2, 10** (45 mg, 0.14 mmol) and *O*-(7-azabenzotriazol-1-yl)- N,N,N',N' tetramethyluronium hexafluorophosphate (HATU) (60 mg, 0.16 mmol) in DMF (1.5 mL), and the mixture was added to a TentaGel- NH_2 resin (100 mg, 0.043 mmol). The resin was shaken at room temperature overnight, washed with DMF, MeOH and CH_2Cl_2 , and dried under vacuum to give resin **Figure 2, 11**. A solution of LiBr (24 mg, 0.28 mmol) in butane-2-one (1.5 mL) was added to the above resin **Figure 2, 11**, and the resin was shaken at 70°C for 72 h. The beads were washed with butane-2-one, DMF and CH_2Cl_2 , and dried to give the resin-bound perfluorosulfonic acid **Figure 2, 12**.

Example 3

This example illustrates the use of perfluorosulfonyl fluoride linker in deoxygenation of various phenols (**Figure 3**).

A mixture of phenol (0.68 mmol), K_2CO_3 (100 mg, 0.72 mmol), resin-bound linker **Figure 3, Structure 4** (80 mg, 0.034 mmol) and DMF (1.0 mL) was shaken at room temperature overnight. The resin was washed with water, DMF and CH_2Cl_2 , and was dried under vacuum overnight to give resin-bound phenol **Figure 3, Structure 13**. To the dry resin **Figure 3, Structure 13** were added $\text{Pd}(\text{OAc})_2$ (6.0 mg, mmol), 1,3-bis(diphenylphosphino)propane (dppp, 16.0 mg, mmol), DMF (1.2-1.4 mL) and a mixture of HCO_2H

(180 μ L) and Et_3N (460 μ L). The mixture was shaken at 85 $^\circ\text{C}$ for 120 min. The polymer beads were filtered and washed with Et_2O . The combined organic phase was washed with aqueous Na_2CO_3 and water, and evaporated to dryness. The residue was dissolved in Et_2O and eluted through a short column of SiO_2 to remove inorganic residues. The crude products were purified by preparative TLC to give the desired products **Figure 3**, **Structure 14a-l** in $> 98\%$ purity.

Example 4

This example illustrates the use of the perfluorosulfonyl halide linkers of the invention for the cleavage/cross coupling of polymer-bound phenols using Suzuki coupling reactions (**Figure 4**).

The resin-bound phenols **Figure 4**, **Structure 15** were prepared as described in Example 1. A mixture of the dry resin **Figure 4**, **Structure 15** (200 mg, 0.07 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (7.2 mg), boronic acid (0.26 mmol) and Et_3N (88 μ L, 0.62 mmol) in DMF (1.5-2.0 mL) was placed in a vial and degassed by blowing N_2 with stirring. The vial was sealed and magnetically stirred at 90 $^\circ\text{C}$ for 8 h. The polymer beads were filtered and washed with Et_2O . The combined organic phase was washed with aqueous Na_2CO_3 solution and water, and evaporated under reduced pressure to dryness. The residue was dissolved in Et_2O and eluted through a short bed of SiO_2 to remove inorganic residues. The crude products were purified by preparative TLC to give the desired products **Figure 4**, **Structure 16a-h** in $> 98\%$ purity.

Example 5

This example illustrates the application of the traceless linker technology to the preparation of a library of 10 biaryl compounds, which further illustrate the scope and generality of the linkers (**Figure 5**).

The resin-bound aldehyde **Figure 5**, **Structure 17** was prepared as described in Example 1. To the dry resin **Figure 5**, **Structure 17** (100 mg, 0.043 mmol) were added amine (0.40 mmol), THF (800 μ L), $\text{Na}(\text{CN})\text{BH}_3$ (1.0 mL) and acetic acid (23 μ L). The mixture was shaken at room temperature overnight. The beads were washed with water, DMF, CH_2Cl_2 and anhydrous THF, and dried under vacuum overnight to give amine resin **Figure 5**, **Structure 18**. To the dry amine resin **Figure 5**, **Structure 18** (100 mg, 0.043

mmol) were added CH_2Cl_2 (1.2-1.6 mL), acid chloride (0.42 mmol), and the resin was cooled in a dry-ice container for 10 min. To the resin was slowly added diisopropylethylamine (88 μL , 0.5 mmol), and the resin was shaken at room temperature overnight. The beads were washed with water, DMF, CH_2Cl_2 , and dried under vacuum overnight to give amide resin **Figure 5, Structure 19**.

A mixture of the dry resin **Figure 5, Structure 19** (200 mg, 0.07 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (7.2 mg), boronic acids (0.26 mmol) and Et_3N (88 μL) in DMF (1.0~1.1 mL) was placed in a vial and degassed by blowing N_2 with stirring. The vial was sealed and magnetically stirred at 90 °C for 8 h. The polymer beads were filtered and washed with Et_2O . The combined organic phase was washed with aqueous Na_2CO_3 solution and water, and evaporated to dryness. The residue was dissolved in Et_2O and eluted through a short bed of SiO_2 to remove inorganic residues. The crude products were purified by preparative TLC to give the desired products **Figure 5, Structure 20a-i** in > 98 % purity.

Example 6

This example illustrates the application of perfluorosulfonyl fluoride linker to a multi-step synthesis of the therapeutic agent, meclizine (**Figure 6**).

A mixture of 3-methyl-4-hydroxybenzaldehyde (100 mg, 0.72 mmol), K_2CO_3 (100 mg, 0.72 mmol), resin-bound linker **Figure 6, Structure 4** (100 mg, 0.043 mmol) and DMF (1.1 mL) was shaken at room temperature overnight. The beads were washed with water, DMF and CH_2Cl_2 , and dried under vacuum overnight to give resin **Figure 6, Structure 21**. To the dry resin **Figure 6, Structure 21** were added 1-(4-chlorobenzhydryl)piperazine (128 mg, 0.40 mmol), THF (800 μL), $\text{Na}(\text{CN})\text{BH}_3$ (1.0 mL) and acetic acid (23 μL). The mixture was shaken at room temperature overnight. The beads were washed with water, DMF and CH_2Cl_2 , and dried under vacuum overnight to give resin **Figure 6, Structure 22**.

To the dry resin **Figure 6, Structure 22** was added $\text{Pd}(\text{OAc})_2$ (8.0 mg, mmol), 1,3-bis(diphenyl-phosphino)propane (dppp, 17.0 mg, mmol), DMF (1.4 mL) and a mixture of HCO_2H (200 μL) and Et_3N (800 μL). The mixture was shaken at 85 °C for 120 min. The polymer beads were filtered and washed with Et_2O . The combined organic phase was washed with aqueous Na_2CO_3 solution and water, and evaporated to dryness. The residue was dissolved in Et_2O and eluted through a short column of Al_2O_3 to remove inorganic

residues. The crude products were purified by preparative TLC to give the desired products **Figure 6, Structure 23** in > 98 % purity. Analytical data of **Figure 6, Structure 23** is identical to that of authentic sample.

5

Example 7

This example illustrates the preparation of the resin-bound perfluorosulfonic acid (**Figure 7**).

10 A mixture of resin-bound linker **Figure 7, Structure 4** (500 mg, 0.15 mmol), THF (1.0 mL) and 0.6 M LiOH in water (1.2 mL) was shaken at room temperature for 2h. The beads were washed with water and CH₂Cl₂, and dried under vacuum overnight to give resin **Figure 7, Structure 24**. ¹⁹F NMR (CDCl₃, δ): -8.00, -43.44.

To the resin **Figure 6, Structure 24** was added 6 M HCl in water (2.0 mL). The mixture was shaken at room temperature for 6 h. The beads were filtered and washed with CH₂Cl₂, 20% TFA in CH₂Cl₂ and CH₂Cl₂, and dried under vacuum overnight to give resin
15 **Figure 7, Structure 25**. ¹⁹F NMR (CDCl₃, δ): -8.00, -43.44, -6.84, -7.41, -12.55, -42.36, -42.88.

Example 8

This example illustrates the use of the resin bound perfluorosulfonic acid as a scavenger.

20 To a solution of benzylamine (0.125 mmol) in 5 mL of THF is added diisopropylethyl amine (0.15 mmol) and benzoylchloride (0.10 mmol). The reaction mixture is stirred at room temperature for 1 hour or until TLC analysis shows consumption of the benzoyl chloride. To this mixture is added perfluorosulfonic acid resin prepared in example 8 (3g, 0.3 mmol/g loading) and the slurry is gently agitated for 1 hour. The slurry
25 is filtered and the resin washed with dichloromethane. The combined filtrates are combined and the solvent removed under reduced pressure to afford N-benzylbenzamide as a solid.

Example 9

This example illustrates the use of the perfluorosulfonyl siloxy resin as a silylating resin.


5 The free acid resin from example 8 (1 g) is washed with dichloromethane. The resin is treated with a solution of TMS-Cl (10 mmol) in 5 mL of dichloromethane and one drop of concentrated sulfuric acid is added. After being gently agitated for 14 h, the resin is washed with dichloromethane and dried under vacuum.

10 To a solution of phenol (0.1 mmol) in dichloromethane is added the TMS-resin from above and the resultant slurry is gently agitated for 1h. The resin is filtered away and the filtrate is either directly submitted for gas chromatographic analysis or the solvent is removed under reduced pressure to yield phenoxytrimethylsilane.

15 All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the
20 appended claims.

WHAT IS CLAIMED IS:

- 1 1. A support-bound activator having the formula:



2
3 wherein

4 L is a linking group component;

5 X is a member selected from the group consisting of F, Cl, OH, and trisubstituted
6 silyloxy;

7 wherein the support-bound activator is covalently attached to a solid or semi-solid
8 support.


1 2. The support-bound activator of claim 1, wherein X is F.

1 3. The support-bound activator of claim 1, wherein X is Cl.

1 4. The support-bound activator of claim 1, wherein X is OH.

1 5. The support-bound activator of claim 1, wherein X is trisubstituted silyloxy.

2 6. The support-bound activator of claim 1, wherein L comprises an activator
3 enhancing portion selected from the group consisting of:



4 and $\xi - Y - Z - (CH_2)_j - (CF_2CF_2)_k - O - (CF_2)_m - \xi$

5 wherein

6 Y is a member selected from the group consisting of a chemical bond, O, C(O), S,
7 and NR¹;

8 Z is a member selected from the group consisting of a chemical bond and C(O);


- 9 R^1 is a member selected from the group consisting of H and (C_1-C_8) alkyl;
 10 each R^2 is independently a member selected from the group consisting of
 11 hydrogen, halogen, (C_1-C_8) alkyl, (C_1-C_8) alkoxy, (C_2-C_8) heteroalkyl, $(C_1-$
 12 $C_8)$ alkylthio, (C_1-C_8) alkylamino, di (C_1-C_8) alkylamino, cyano, nitro, and
 13 (C_1-C_8) alkylsulfonyl;
 14 the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
 15 7, 8, 9, and 10;
 16 the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and
 17 6;
 18 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
 19 7, 8, 9, and 10;
 20 the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;
 21 the subscript m is an integer selected from the group consisting of 2, and 3; and
 22 the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.

1 7. The support-bound activator of claim 1, wherein said solid or semi-solid
 2 support is a particulate support.

1 8. The support-bound activator of claim 1, wherein said solid or semi-solid
 2 support comprises a material selected from the group consisting of polystyrene, controlled
 3 pore glass, polyacrylamide, poly(ethyleneglycol)monomethyl ether, polyethylene glycol,
 4 silica gel, cellulose, acrylic acid grafted polypropylene, polystyrene modified by
 5 polyethylene glycol, and combinations thereof.

1 9. The support-bound activator of claim 1, wherein said solid or semi-solid
 2 support comprises polystyrene modified by polyethylene glycol

1 10. An activated support comprising a solid or semi-solid support; and at least one
 2 support-bound activator having the formula:
 3





5 wherein

L is a linking group component;

X is a member selected from the group consisting of F, Cl, OH, and trisubstituted silyloxy; and

wherein the support-bound activator is covalently attached to the solid or semi-solid support.

11. The activated support of claim 10, wherein L comprises an activator enhancing portion selected from the group consisting of:



wherein

Y is a member selected from the group consisting of a chemical bond, O, C(O), S, and NR¹;

Z is a member selected from the group consisting of a chemical bond and C(O);

R¹ is a member selected from the group consisting of H and (C₁-C₈)alkyl;

each R² is independently a member selected from the group consisting of

hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro, and (C₁-C₈)alkylsulfonyl;

the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,

7, 8, 9, and 10;

the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and

6;

the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,

7, 8, 9, and 10;

the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;

the subscript m is an integer selected from the group consisting of 2, and 3; and

the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.


12. The activated support of claim 10, wherein said solid or semi-solid support is a particulate support.

13. The activated support of claim 10, wherein said solid or semi-solid support comprises a material selected from the group consisting of polystyrene, controlled pore glass, polyacrylamide, poly(ethyleneglycol)monomethyl ether, polyethylene glycol, silica gel, cellulose, acrylic acid grafted polypropylene, polystyrene modified by polyethylene glycol, and combinations thereof.

14. The activated support of claim 10, wherein said solid or semi-solid support comprises polystyrene modified by polyethylene glycol.

15. A kit comprising a predetermined amount of the activated support of claim 10




16. A support-activated target comprising a solid or semi-solid support; an activating group covalently attached to the solid or semi-solid support, wherein the activating group has the formula:



and wherein L is a linking group component;

and a target group covalently attached to the activating group; wherein the target group can be cleaved from the activating group by a nucleophile.

17. The support-activated target of claim 16, wherein L comprises an activator enhancing portion selected from the group consisting of:



wherein

5 Y is a member selected from the group consisting of a chemical bond, O, C(O), S,
6 and NR¹;

7 Z is a member selected from the group consisting of a chemical bond and C(O);

8 R¹ is a member selected from the group consisting of H and (C₁-C₈)alkyl;

9 each R² is independently a member selected from the group consisting of
10 hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-
11 C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro, and
12 (C₁-C₈)alkylsulfonyl;

13 the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
14 7, 8, 9, and 10;

15 the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and
16 6;

17 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
18 7, 8, 9, and 10;

19 the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;

20 the subscript m is an integer selected from the group consisting of 2, and 3; and

21 the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.


1 18. The support-activated target of claim 16, wherein said solid or semi-solid
2 support is a particulate support.

1 19. The support-activated target of claim 16, wherein said solid or semi-solid
2 support is composed of a material selected from the group consisting of polystyrene,
3 controlled pore glass, polyacrylamide, poly(ethyleneglycol)monomethyl ether,
4 polyethylene glycol, silica gel, cellulose, acrylic acid grafted polypropylene, polystyrene
5 modified by polyethylene glycol, and combinations thereof.

1 20. The support-activated target of claim 16, wherein said solid or semi-solid
2 support comprises polystyrene modified by polyethylene glycol.



1 21. A library of support-activated targets comprising a plurality of support-
2 activated target members, wherein each support-activated target member further comprises

3 a solid or semi-solid support; an activating group covalently attached to the solid or semi-
 4 solid support, wherein the activating group has the formula:



5
 6
 7 wherein L is an linking group component; and a target group covalently attached to the
 8 activating group; wherein the target group of at least one support-activated target member
 9 in the library is different from the target group of at least one other support-activated target
 10 member in the library.

1 22. The library of support-activated targets of claim 21, wherein L comprises an
 2 activator enhancing portion selected from the group consisting of:



3
 4 wherein

5 Y is a member selected from the group consisting of a chemical bond, O, C(O), S,
 6 and NR¹;

7 Z is a member selected from the group consisting of a chemical bond and C(O);

8 R¹ is a member selected from the group consisting of H and (C₁-C₈)alkyl;

9 each R² is independently a member selected from the group consisting of

10 hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-
 11 C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro, and
 12 (C₁-C₈)alkylsulfonyl;

13 the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
 14 7, 8, 9, and 10;


15 the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and
 16 6;
 17 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
 18 7, 8, 9, and 10;
 19 the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;
 20 the subscript m is an integer selected from the group consisting of 2, and 3; and
 21 the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.

1 23. The library of support-activated targets of claim 21, wherein said solid or semi-
 2 solid support is a particulate support.

1 24. The library of support-activated targets of claim 21, wherein said solid or semi-
 2 solid support is composed of a material selected from the group consisting of polystyrene,
 3 controlled pore glass, polyacrylamide, poly(ethyleneglycol)monomethyl ether,
 4 polyethylene glycol, silica gel, cellulose, acrylic acid grafted polypropylene, polystyrene
 5 modified by polyethylene glycol, and combinations thereof.

1 25. The library of support-activated targets of claim 21, wherein said solid or semi-
 2 solid support comprises polystyrene modified by polyethylene glycol.

1 26. A strongly acidic support comprising a solid or semi-solid support; and at least
 2 one support-bound strongly acidic group having the formula:





3
 4 wherein

5 L is a linking group component; and

6 X is OH;

7 wherein the support-bound strongly acidic group is covalently attached to the solid
 8 or semi-solid support.

1 27. The strongly acidic support of claim 26, wherein L comprises an activator
 2 enhancing portion selected from the group consisting of:



3

4 wherein

5 Y is a member selected from the group consisting of a chemical bond, O, C(O), S,
6 and NR¹;

7 Z is a member selected from the group consisting of a chemical bond and C(O);

8 R¹ is a member selected from the group consisting of H and (C₁-C₈)alkyl;

9 each R² is independently a member selected from the group consisting of

10 hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-
11 C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro, and
12 (C₁-C₈)alkylsulfonyl;

13 the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
14 7, 8, 9, and 10;

15 the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and
16 6;

17 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
18 7, 8, 9, and 10;

19 the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;

20 the subscript m is an integer selected from the group consisting of 2, and 3; and


21 the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.

1 28. The strongly acidic support of claim 26, wherein said solid or semi-solid
2 support is a particulate support.

1 29. The strongly acidic support of claim 26, wherein said solid or semi-solid
2 support is composed of a material selected from the group consisting of polystyrene,
3 controlled pore glass, polyacrylamide, poly(ethyleneglycol)monomethyl ether,
4 polyethylene glycol, silica gel, cellulose, acrylic acid grafted polypropylene, polystyrene
5 modified by polyethylene glycol, and combinations thereof.

1 30. The strongly acidic support of claim 26, wherein said solid or semi-solid
2 support comprises polystyrene modified by polyethylene glycol.

1 31. A silylating support comprising a solid or semi-solid support; and at least one
2 support-bound silylating group having the formula:




3
4 wherein

5 L is a linking group component; and

6 X is a trisubstituted silyloxy;

7 wherein the support-bound silylating group is covalently attached to the solid or
8 semi-solid support.

1 32. The silylating support of claim 31, wherein L comprises an activator enhancing
2 portion selected from the group consisting of:



3 and $\xi - Y - Z - (CH_2)_j - (CF_2CF_2)_k - O - (CF_2)_m - \xi$

4 wherein

5 Y is a member selected from the group consisting of a chemical bond, O, C(O), S,
6 and NR¹;

7 Z is a member selected from the group consisting of a chemical bond and C(O);

8 R¹ is a member selected from the group consisting of H and (C₁-C₈)alkyl;

9 each R² is independently a member selected from the group consisting of

10 hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-
11 C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro, and
12 (C₁-C₈)alkylsulfonyl;

13 the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,

14 7, 8, 9, and 10;

15 the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and
16 6;
17 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
18 7, 8, 9, and 10;
19 the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;
20 the subscript m is an integer selected from the group consisting of 2, and 3; and
21 the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.

1 33. The silylating support of claim 31, wherein said solid or semi-solid support is a
2 particulate support.

1 34. The silylating support of claim 31, wherein said solid or semi-solid support is
2 composed of a material selected from the group consisting of polystyrene, controlled pore
3 glass, polyacrylamide, poly(ethyleneglycol)monomethyl ether, polyethylene glycol, silica
4 gel, cellulose, acrylic acid grafted polypropylene, polystyrene modified by polyethylene
5 glycol, and combinations thereof.

1 35. The silylating support of claim 31, wherein said solid or semi-solid support
2 comprises polystyrene modified by polyethylene glycol.

1 36. The silylating support of claim 31, wherein X is a tri(C1-C8) silyloxy.

1 37. The silylating support of claim 31, wherein X is a member selected from the
2 group consisting of $\text{OSi}(\text{CH}_3)_3$ and $\text{OSi}(\text{CH}_3)_2\text{t-Bu}$.

1 38. A kit for silylating a target comprising a predetermined amount of the silylating
2 support of claim 31.

3 39. A method for covalently attaching a nucleophile to a compound having a
4 hydroxy group or an enolizable ketone, said method comprising,

5 (a) contacting said compound with a support-bound activator of claim 1, to form
6 an activated complex; and

7 (b) contacting said activated complex with a reagent, said reagent comprising a
8 nucleophile; wherein said contacting comprises conditions sufficient to covalently attach
9 said nucleophile to said compound.

1 40. A method in accordance with claim 39 wherein said conditions sufficient to
2 covalently attach said nucleophile to said compound includes addition of a transition metal
3 catalyst.

1 41. A method in accordance with claim 39 wherein said reagent is a member
2 selected from the group consisting of an organostannane compound, an organolithium
3 compound, an organozinc compound, an organoboron compound, an organoaluminum
4 compound, a Grignard reagent, an organosilicon compound, an organocopper compound,
5 a thiol, a dialkylphosphite, an amine, a metal halide, and a halogen.

1 42. A method in accordance with claim 39, wherein said nucleophile is a member
2 selected from the group consisting of an amine, a halogen anion, an aryl moiety, an alkyl
3 moiety, and a hydride.


1 43. A method in accordance with claim 39, wherein said nucleophile is hydride.

1 44. A method in accordance with claim 39, wherein said reagent is an aryl boronic
2 acid.

1 45. A method in accordance with claim 39, wherein said nucleophile is an ^{18}F
2 anion.

1 46. A method in accordance with claim 39, wherein said nucleophile is an $^{11}\text{CH}_3$
2 anion.

1 47. A linker reagent having the formula:



2
3 wherein

4 L is a linking group component; and

5 A is an attaching group.


1 48. A linker reagent of claim 47, wherein said attaching group is a member
2 selected from the group consisting of NH_2 , NHR , CO_2H , CO_2R , COCl , OH , SH , and

3 protected forms thereof, wherein each R is independently a member selected from the
 4 group consisting of substituted (C₁-C₈)alkyl, unsubstituted (C₁-C₈)alkyl, substituted aryl,
 5 and unsubstituted aryl.

1 49. A linker reagent of claim 47, wherein said attaching group is CO₂H.

1 50. A linker reagent of claim 47, wherein said attaching group is COCl.

1 51. A linker reagent of claim 47, wherein L is a member selected from the group
 2 consisting of:



3 and $\xi-Y-Z-(CH_2)_j-(CF_2CF_2)_k-O-(CF_2)_m-\xi$

4 wherein

5 Y is a member selected from the group consisting of a chemical bond, O, C(O), S,
 6 and NR¹;

7 Z is a member selected from the group consisting of a chemical bond and C(O);

8 R¹ is a member selected from the group consisting of H and (C₁-C₈)alkyl;

9 each R² is independently a member selected from the group consisting of

10 hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)alkoxy, (C₂-C₈)heteroalkyl, (C₁-
 11 C₈)alkylthio, (C₁-C₈)alkylamino, di(C₁-C₈)alkylamino, cyano, nitro, and
 12 (C₁-C₈)alkylsulfonyl;

13 the subscript g is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
 14 7, 8, 9, and 10;

15 the subscript i is an integer selected from the group consisting of 1, 2, 3, 4, 5, and
 16 6;


17 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
 18 7, 8, 9, and 10;

19 the subscript k is an integer selected from the group consisting of 1, 2, 3, and 4;

20 the subscript m is an integer selected from the group consisting of 2, and 3; and

21 the subscript n is an integer selected from the group consisting of 0, 1, 2, 3, and 4.

1 52. A linker reagent in accordance with claim 47, having the formula:



3 wherein

4 the subscript j is an integer selected from the group consisting of 0, 1, 2, 3,
5 4, 5, 6, 7, 8, 9, and 10; and

6 the subscript k is an integer selected from the group consisting of 1, 2, 3,
7 and 4.

1 53. A linker reagent in accordance with claim 47, wherein the subscript j is 1 and
2 the subscript k is 1.

1 54. A linker reagent in accordance with claim 47, having the formula:



3 wherein the subscript g is an integer selected from the group consisting of 3, 4, 5,
4 and 6; and

5 the subscript i is an integer selected from the group consisting of 0, 1, 2, 3, 4, 5, 6,
6 7, 8, 9, and 10.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 July 2002 (18.07.2002)

PCT

(10) International Publication Number
WO 02/055026 A3

(51) International Patent Classification⁷: **G01N 33/543**

(21) International Application Number: **PCT/US02/00839**

(22) International Filing Date: **11 January 2002 (11.01.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/261,525 **12 January 2001 (12.01.2001)** **US**

(71) Applicant (for all designated States except US): **AFFY-MAX, INC.** [US/US]; 4001 Miranda Avenue, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HOLMES, Christopher** [US/US]; 13633 Westover Drive, Saratoga, CA 95070 (US). **PAN, Yijun** [CN/US]; 3791 Boswell Terrace, Fremont, CA 94536 (US).

(74) Agents: **HOSTETLER, Michael, J.** et al.; Brobeck, Phleger & Harrison, LLP, 12390 El Camino Real, San Diego, CA 92130 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:
14 November 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PERFLUORO SULFONYL HALIDES AND RELATED SPECIES AS POLYMER SUPPORT MODIFIERS**



(I)

(57) Abstract: Activated Supports, support-bound activators, strongly acidic supports, and silylating supports are provided; the activated support having the formula (I) wherein L is a linking group component; X is F, CL, OH, and trisubstituted silyloxy; and the shaded circle represents a solid or semi-solid support. Methods of using the activated supports in solid phase organic synthesis are also provided.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/00889

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : G01N 33/543

US Cl. : 568/32, 33; 436/518; 435/DIG40, DIG 42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 568/32, 33; 436/518; 435/DIG40, DIG 42

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAPLUS, WEST, STIC

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AKHTAR et al. Solid-phase synthesis on functionalized fluorophore resins. Part I: Nafion resin sulfonamide-immobilised carboxylic acid derivatives and aryl vinyl sulfones. Tet. Lett., June 2000, Vol. 41, pp. 4487-4491, see entire document	1-38, 47-54
A	SMITH, E.M. A Polymer-Supported Silyl Triflate and Subsequent Functionalization: Synthesis and Solid-Phase Diels-Alder Reactions of Silyloxydienes. Tet. Lett., April 1999, Vol. 40, pp. 3285-3288, see entire document	1-54
A	HU et al. Ester Enolate Claisen Rearrangement Using a Polymer-Supported Silyl Triflate. Tet. Lett., April 1999, Vol. 40, pp. 3289-3292, see entire document	1-38, 47-54

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"G" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

24 MAY 2002

Date of mailing of the international search report

12 JUL 2002

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer:

STEVEN C. TIZIO

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/00839

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ZARAGOZA, Florencio New Sulfur- and Selenium-Based Traceless Linkers - More than just Linkers? Angew. Chem. Int. Ed., June 2000, Vol. 39, No. 12, pp. 2077-2079, see entire document	47-54
A	US 5,919,523 A (SUNDBERG et al.) 06 July 1999, see entire document, especially column 2.	1-54

Form PCT/ISA/210 (continuation of second sheet) (July 1998)★

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US02/00889

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-2, 26-38, drawn to the support-bound activator, support-bound target, and strongly acidic support.

Group II, claim(s) 21-25, drawn to the library of support-activated targets.

Group III, claim(s) 39-46, drawn to the method for covalently attaching a nucleophile to a compound having a hydroxy group or an enolizable ketone.

Group IV, claim(s) 47-54, drawn to the linker reagent.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The library of support-activated targets of Group II can have more than one target compared to the invention of Group I. The method of Group III relates to a method for a chemical reaction. Finally, the linker reagent of Group IV has a different chemical formula than the compounds in Groups I and II. In addition, the linker reagent in Group IV is not attached to a solid support.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

The claims are deemed to correspond to the species listed above in the following manner:

1. Applicant is requested to elect either a halogen or trisubstituted silyloxy for X.
2. Applicant is requested to elect a single species of activator enhancing portion.

The following claims are generic: 1-54

Each of the species are distinct from each other because the compounds are structurally and functionally different from each other and do not require the other for ultimate use; the species election for examination purposes as indicated is proper.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.